

EWSR1 Blocking Peptide (Center)

Synthetic peptide

Catalog # BP20981a

Specification**EWSR1 Blocking Peptide (Center) - Product Information**Primary Accession [Q01844](#)Other Accession [Q61545](#)**EWSR1 Blocking Peptide (Center) - Additional Information****Gene ID** 2130**Other Names**RNA-binding protein EWS, EWS oncogene,
Ewing sarcoma breakpoint region 1 protein,
EWSR1, EWS**Target/Specificity**The synthetic peptide sequence is selected
from aa 328-342 of HUMAN EWSR1**Format**Peptides are lyophilized in a solid powder
format. Peptides can be reconstituted in
solution using the appropriate buffer as
needed.**Storage**Maintain refrigerated at 2-8°C for up to 6
months. For long term storage store at
-20°C.**Precautions**This product is for research use only. Not
for use in diagnostic or therapeutic
procedures.**EWSR1 Blocking Peptide (Center) - Protein Information****Name** EWSR1**Synonyms** EWS**Function**Might normally function as a transcriptional
repressor. EWS- fusion-proteins (EFPS) may**EWSR1 Blocking Peptide (Center) - Background**

Might normally function as a transcriptional repressor. EWS-fusion-proteins (EFPS) may play a role in the tumorigenic process. They may disturb gene expression by mimicking, or interfering with the normal function of CTD-POLII within the transcription initiation complex. They may also contribute to an aberrant activation of the fusion protein target genes.

EWSR1 Blocking Peptide (Center) - References

Delattre O.,et al.Nature 359:162-165(1992).
Plougastel B.,et al.Genomics
18:609-615(1993).
Zucman-Rossi J.,et al.Submitted (MAY-1998) to
the EMBL/GenBank/DDBJ databases.
Collins J.E.,et al.Genome Biol.
5:R84.1-R84.11(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).

play a role in the tumorigenic process. They may disturb gene expression by mimicking, or interfering with the normal function of CTD-POLII within the transcription initiation complex. They may also contribute to an aberrant activation of the fusion protein target genes.

Cellular Location

Nucleus. Cytoplasm. Cell membrane.

Note=Relocates from cytoplasm to ribosomes upon PTK2B/FAK2 activation

Tissue Location

Ubiquitous.

**EWSR1 Blocking Peptide (Center) -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)