

SNAP25 Blocking Peptide (S187)

Synthetic peptide

Catalog # BP21014a

Specification**SNAP25 Blocking Peptide (S187) - Product Information**

Primary Accession [P60880](#)
Other Accession [P60881](#), [P60879](#),
[P60878](#), [Q170Q3](#),
[Q5TZ66](#)

SNAP25 Blocking Peptide (S187) - Additional Information

Gene ID 6616

Other Names

Synaptosomal-associated protein 25,
SNAP-25, Super protein, SUP,
Synaptosomal-associated 25 kDa protein,
SNAP25, SNAP

Target/Specificity

The synthetic peptide sequence is selected
from aa 179-193 of HUMAN SNAP25

Format

Peptides are lyophilized in a solid powder
format. Peptides can be reconstituted in
solution using the appropriate buffer as
needed.

Storage

Maintain refrigerated at 2-8°C for up to 6
months. For long term storage store at
-20°C.

Precautions

This product is for research use only. Not
for use in diagnostic or therapeutic
procedures.

SNAP25 Blocking Peptide (S187) - Protein Information

Name SNAP25

Synonyms SNAP

SNAP25 Blocking Peptide (S187) - Background

t-SNARE involved in the molecular regulation
of neurotransmitter release. May play an
important role in the synaptic function of
specific neuronal systems. Associates with
proteins involved in vesicle docking and
membrane fusion. Regulates plasma
membrane recycling through its interaction
with CENPF.

SNAP25 Blocking Peptide (S187) - References

Bark I.C.,et al.Gene 139:291-292(1994).
Zhao N.,et al.Gene 145:313-314(1994).
Jagadish M.N.,et al.Biochem. J.
317:945-954(1996).
Kalnine N.,et al.Submitted (OCT-2004) to the
EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Function

t-SNARE involved in the molecular regulation of neurotransmitter release. May play an important role in the synaptic function of specific neuronal systems. Associates with proteins involved in vesicle docking and membrane fusion. Regulates plasma membrane recycling through its interaction with CENPF. Modulates the gating characteristics of the delayed rectifier voltage-dependent potassium channel KCNB1 in pancreatic beta cells.

Cellular Location

Cytoplasm, perinuclear region
{ECO:0000250|UniProtKB:P60879}. Cell membrane
{ECO:0000250|UniProtKB:P60881};
Lipid-anchor
{ECO:0000250|UniProtKB:P60879}. Cell junction, synapse, synaptosome
{ECO:0000250|UniProtKB:P60879}. Photoreceptor inner segment
{ECO:0000250|UniProtKB:P60879}.
Note=Membrane association requires palmitoylation. Expressed throughout cytoplasm, concentrating at the perinuclear region. Colocalizes with KCNB1 at the cell membrane (By similarity). Colocalizes with PLCL1 at the cell membrane (By similarity).
{ECO:0000250|UniProtKB:P60879, ECO:0000250|UniProtKB:P60881}

Tissue Location

Neurons of the neocortex, hippocampus, piriform cortex, anterior thalamic nuclei, pontine nuclei, and granule cells of the cerebellum

**SNAP25 Blocking Peptide (S187) -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)