

Mouse Ptk7 Blocking Peptide (C-term)
Synthetic peptide
Catalog # BP21199b**Specification****Mouse Ptk7 Blocking Peptide (C-term) - Product Information**Primary Accession [Q8BKG3](#)**Mouse Ptk7 Blocking Peptide (C-term) - Additional Information****Gene ID** 71461**Other Names**

Inactive tyrosine-protein kinase 7, Protein chuzhoi, Protein-tyrosine kinase 7, Pseudo tyrosine kinase receptor 7, Tyrosine-protein kinase-like 7, Ptk7

Target/Specificity

The synthetic peptide sequence is selected from aa 722-737 of HUMAN Ptk7

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Mouse Ptk7 Blocking Peptide (C-term) - Protein Information**Name** Ptk7**Function**

Inactive tyrosine kinase involved in Wnt signaling pathway. Component of both the non-canonical (also known as the Wnt/planar cell polarity signaling) and the

Mouse Ptk7 Blocking Peptide (C-term) - Background

Inactive tyrosine kinase involved in Wnt signaling pathway. Component of both the non-canonical (also known as the Wnt/planar cell polarity signaling) and the canonical Wnt signaling pathway. Functions in cell adhesion, cell migration, cell polarity, proliferation, actin cytoskeleton reorganization and apoptosis. Has a role in embryogenesis, epithelial tissue organization and angiogenesis.

Mouse Ptk7 Blocking Peptide (C-term) - References

Jung J.-W., et al. Gene 328:75-84(2004).
Carninci P., et al. Science 309:1559-1563(2005).
Daigo Y., et al. Submitted (FEB-2001) to the EMBL/GenBank/DDBJ databases.
Lu X., et al. Nature 430:93-98(2004).
Gundry R.L., et al. Mol. Cell. Proteomics 8:2555-2569(2009).

canonical Wnt signaling pathway. Functions in cell adhesion, cell migration, cell polarity, proliferation, actin cytoskeleton reorganization and apoptosis. Has a role in embryogenesis, epithelial tissue organization and angiogenesis.

Cellular Location

Membrane; Single-pass type I membrane protein. Cell junction. Note=Colocalizes with MMP14 at cell junctions. Also localizes at the leading edge of migrating cells

Tissue Location

Expressed at high levels in lung and un-pregnant uterus among adult tissues, and in the tail, limbs, somites, gut and craniofacial regions among embryonic tissues

Mouse Ptk7 Blocking Peptide (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)