

**Mouse Kdm6a Blocking Peptide (C-term)**  
**Synthetic peptide**  
**Catalog # BP21225b****Specification****Mouse Kdm6a Blocking Peptide (C-term) - Product Information**Primary Accession [O70546](#)**Mouse Kdm6a Blocking Peptide (C-term) - Additional Information****Gene ID** 22289**Other Names**

Lysine-specific demethylase 6A, 11411-, Histone demethylase UTX, Ubiquitously transcribed TPR protein on the X chromosome, Ubiquitously transcribed X chromosome tetratricopeptide repeat protein, Kdm6a, Utx

**Target/Specificity**

The synthetic peptide sequence is selected from aa 1041-1054 of HUMAN Kdm6a

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Mouse Kdm6a Blocking Peptide (C-term) - Protein Information****Name** Kdm6a**Synonyms** Utx**Function****Mouse Kdm6a Blocking Peptide (C-term) - Background**

Histone demethylase that specifically demethylates 'Lys- 27' of histone H3, thereby playing a central role in histone code. Demethylates trimethylated and dimethylated but not monomethylated H3 'Lys-27'. Plays a central role in regulation of posterior development, by regulating HOX gene expression. Demethylation of 'Lys-27' of histone H3 is concomitant with methylation of 'Lys-4' of histone H3, and regulates the recruitment of the PRC1 complex and monoubiquitination of histone H2A (By similarity).

**Mouse Kdm6a Blocking Peptide (C-term) - References**

Church D.M.,et al.PLoS Biol. 7:E1000112-E1000112(2009).  
Greenfield A.,et al.Hum. Mol. Genet. 7:737-742(1998).  
Grbavec D.,et al.Biochem. J. 337:13-17(1999).  
Ballif B.A.,et al.Mol. Cell. Proteomics 3:1093-1101(2004).  
Wang A.H.,et al.EMBO J. 32:1075-1086(2013).

Histone demethylase that specifically demethylates 'Lys-27' of histone H3, thereby playing a central role in histone code. Demethylates trimethylated and dimethylated but not monomethylated H3 'Lys-27'. Plays a central role in regulation of posterior development, by regulating HOX gene expression. Demethylation of 'Lys-27' of histone H3 is concomitant with methylation of 'Lys-4' of histone H3, and regulates the recruitment of the PRC1 complex and monoubiquitination of histone H2A (By similarity). Plays a demethylase-independent role in chromatin remodeling to regulate T-box family member-dependent gene expression (PubMed:<a href="http://www.uniprot.org/citations/21095589" target="\_blank">21095589</a>).

**Cellular Location**

Nucleus.

**Tissue Location**

Expressed in brain, heart and spleen.

**Mouse Kdm6a Blocking Peptide (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)