

ITGA3 Blocking Peptide (C-term)

Synthetic peptide

Catalog # BP21353b

Specification**ITGA3 Blocking Peptide (C-term) - Product Information**Primary Accession [P26006](#)**ITGA3 Blocking Peptide (C-term) - Additional Information****Gene ID** 3675**Other Names**

Integrin alpha-3, CD49 antigen-like family member C, FRP-2, Galactoprotein B3, GAPB3, VLA-3 subunit alpha, CD49c, Integrin alpha-3 heavy chain, Integrin alpha-3 light chain, ITGA3, MSK18

Target/Specificity

The synthetic peptide sequence is selected from aa 875-889 of HUMAN ITGA3

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ITGA3 Blocking Peptide (C-term) - Protein Information**Name** ITGA3**Synonyms** MSK18**Function**

Integrin alpha-3/beta-1 is a receptor for

ITGA3 Blocking Peptide (C-term) - Background

Integrin alpha-3/beta-1 is a receptor for fibronectin, laminin, collagen, epiligrin, thrombospondin and CSPG4. Integrin alpha-3/beta-1 provides a docking site for FAP (seprase) at invadopodia plasma membranes in a collagen-dependent manner and hence may participate in the adhesion, formation of invadopodia and matrix degradation processes, promoting cell invasion. Alpha-3/beta-1 may mediate with LGALS3 the stimulation by CSPG4 of endothelial cells migration.

ITGA3 Blocking Peptide (C-term) - References

Takada Y.,et al.J. Cell Biol. 115:257-266(1991).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Zody M.C.,et al.Nature 440:1045-1049(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Tsuji T.,et al.J. Biochem. 109:659-665(1991).

fibronectin, laminin, collagen, epiligrin, thrombospondin and CSPG4. Integrin alpha-3/beta-1 provides a docking site for FAP (seprase) at invadopodia plasma membranes in a collagen-dependent manner and hence may participate in the adhesion, formation of invadopodia and matrix degradation processes, promoting cell invasion. Alpha-3/beta-1 may mediate with LGALS3 the stimulation by CSPG4 of endothelial cells migration.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell membrane; Lipid-anchor. Cell projection, invadopodium membrane; Single-pass type I membrane protein. Cell projection, filopodium membrane; Single-pass type I membrane protein. Note=Enriched preferentially at invadopodia, cell membrane protrusions that correspond to sites of cell invasion, in a collagen-dependent manner.

Tissue Location

Isoform 1 is widely expressed. Isoform 2 is expressed in brain and heart. In brain, both isoforms are exclusively expressed on vascular smooth muscle cells, whereas in heart isoform 1 is strongly expressed on vascular smooth muscle cells, isoform 2 is detected only on endothelial vein cells.

**ITGA3 Blocking Peptide (C-term) -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)