

SRY Blocking Peptide (Center)
Synthetic peptide
Catalog # BP21539c**Specification****SRY Blocking Peptide (Center) - Product Information**Primary Accession [Q05066](#)**SRY Blocking Peptide (Center) - Additional Information****Gene ID** 6736**Other Names**Sex-determining region Y protein,
Testis-determining factor, SRY, TDF**Target/Specificity**

The synthetic peptide sequence is selected from aa 81-92 of HUMAN SRY

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SRY Blocking Peptide (Center) - Protein Information**Name** SRY{ECO:0000303|PubMed:1695712,
ECO:0000312|HGNC:HGNC:11311}**Function**

Transcriptional regulator that controls a genetic switch in male development (PubMed:<a href="http://www.uniprot.org/citations/11563911"

SRY Blocking Peptide (Center) - Background

Transcriptional regulator that controls a genetic switch in male development. It is necessary and sufficient for initiating male sex determination by directing the development of supporting cell precursors (pre-Sertoli cells) as Sertoli rather than granulosa cells (By similarity). In male adult brain involved in the maintenance of motor functions of dopaminergic neurons (By similarity). Involved in different aspects of gene regulation including promoter activation or repression (By similarity). Promotes DNA bending. SRY HMG box recognizes DNA by partial intercalation in the minor groove. Also involved in pre-mRNA splicing. Binds to the DNA consensus sequence 5'-[AT]AACAA[AT]-3'.

SRY Blocking Peptide (Center) - References

Sinclair A.H.,et al.Nature 346:240-244(1990).
Su H.,et al.Am. J. Hum. Genet. 52:24-38(1993).
Behlke M.A.,et al.Genomics 17:736-739(1993).
Whitfield L.S.,et al.Genomics 27:306-311(1995).
Ferrari S.,et al.EMBO J. 11:4497-4506(1992).

target="_blank">11563911). It is necessary and sufficient for initiating male sex determination by directing the development of supporting cell precursors (pre-Sertoli cells) as Sertoli rather than granulosa cells (PubMed:16996051, PubMed:16414182). Involved in different aspects of gene regulation including promoter activation or repression (PubMed:9525897). Binds to the DNA consensus sequence 5'-[AT]AACAA[AT]-3' (PubMed:1425584, PubMed:8265659, PubMed:8159753, PubMed:11563911, PubMed:15170344). SRY HMG box recognizes DNA by partial intercalation in the minor groove and promotes DNA bending (PubMed:1425584, PubMed:8265659, PubMed:8159753, PubMed:11563911, PubMed:15170344, PubMed:16762365). Also involved in pre-mRNA splicing (PubMed:11818535). In male adult brain involved in the maintenance of motor functions of

dopaminergic neurons (By similarity).

Cellular Location

Nucleus speckle. Cytoplasm. Nucleus

Note=Acetylation contributes to its nuclear localization and deacetylation by HDAC3

induces a cytoplasmic delocalization

(PubMed:15297880). Colocalizes with SOX6

in speckles (PubMed:11818535) Colocalizes with CAML in the nucleus

(PubMed:15746192). Colocalizes in the

nucleus with ZNF208 isoform KRAB-O and

tyrosine hydroxylase (TH) (By similarity).

{ECO:0000250|UniProtKB:Q05738,

ECO:0000269|PubMed:11818535,

ECO:0000269|PubMed:15297880,

ECO:0000269|PubMed:15746192}

SRY Blocking Peptide (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)