



### **TST Blocking Peptide (C-Term)**

Synthetic peptide Catalog # BP21669b

## **Specification**

TST Blocking Peptide (C-Term) - Product Information

Primary Accession <u>Q16762</u>

TST Blocking Peptide (C-Term) - Additional Information

**Gene ID** 7263

#### **Other Names**

Thiosulfate sulfurtransferase, Rhodanese, TST

#### **Target/Specificity**

The synthetic peptide sequence is selected from aa 218-232 of HUMAN TST

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

TST Blocking Peptide (C-Term) - Protein Information

## Name TST

#### **Function**

Formation of iron-sulfur complexes, cyanide detoxification or modification of sulfur-containing enzymes. Other thiol compounds, besides cyanide, can act as sulfur ion acceptors. Also has weak mercaptopyruvate sulfurtransferase (MST)

# TST Blocking Peptide (C-Term) - Background

Formation of iron-sulfur complexes, cyanide detoxification or modification of sulfur-containing enzymes. Other thiol compounds, besides cyanide, can act as sulfur ion acceptors. Also has weak mercaptopyruvate sulfurtransferase (MST) activity (By similarity). Together with MRPL18, acts as a mitochondrial import factor for the cytosolic 5S rRNA. Only the nascent unfolded cytoplasmic form is able to bind to the 5S rRNA.

## TST Blocking Peptide (C-Term) - References

Aita N.,et al.Biochem. Biophys. Res. Commun. 231:56-60(1997). Collins J.E.,et al.Genome Biol. 5:R84.1-R84.11(2004). Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004). Dunham I.,et al.Nature 402:489-495(1999).





Tel: 858.875.1900 Fax: 858.622.0609

activity (By similarity). Together with MRPL18, acts as a mitochondrial import factor for the cytosolic 5S rRNA. Only the nascent unfolded cytoplasmic form is able to bind to the 5S rRNA.

**Cellular Location** Mitochondrion matrix.

## **TST Blocking Peptide (C-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides