

NCOA2 Blocking Peptide (C-Term)
Synthetic peptide
Catalog # BP21793b**Specification****NCOA2 Blocking Peptide (C-Term) - Product Information**Primary Accession [Q15596](#)**NCOA2 Blocking Peptide (C-Term) - Additional Information****Gene ID** 10499**Other Names**

Nuclear receptor coactivator 2, NCoA-2, Class E basic helix-loop-helix protein 75, bHLHe75, Transcriptional intermediary factor 2, hTIF2, NCOA2, BHLHE75, SRC2, TIF2

Target/Specificity

The synthetic peptide sequence is selected from aa 1451-1462 of HUMAN NCOA2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

NCOA2 Blocking Peptide (C-Term) - Protein Information**Name** NCOA2**Synonyms** BHLHE75, SRC2, TIF2**Function**

Transcriptional coactivator for steroid

NCOA2 Blocking Peptide (C-Term) - Background

Transcriptional coactivator for steroid receptors and nuclear receptors. Coactivator of the steroid binding domain (AF-2) but not of the modulating N-terminal domain (AF-1). Required with NCOA1 to control energy balance between white and brown adipose tissues. Critical regulator of glucose metabolism regulation, acts as RORA coactivator to specifically modulate G6PC expression. Involved in the positive regulation of the transcriptional activity of the glucocorticoid receptor NR3C1 by sumoylation enhancer RWDD3. Positively regulates the circadian clock by acting as a transcriptional coactivator for the CLOCK-ARNTL/BMAL1 heterodimer (By similarity).

NCOA2 Blocking Peptide (C-Term) - References

Voegel J.J., et al. EMBO J. 15:3667-3675(1996).
Carapeti M., et al. Blood 91:3127-3133(1998).
Voegel J.J., et al. EMBO J. 17:507-519(1998).
Fryer C.J., et al. Nature 393:88-91(1998).
Atkins G.B., et al. Mol. Endocrinol. 13:1550-1557(1999).

receptors and nuclear receptors. Coactivator of the steroid binding domain (AF-2) but not of the modulating N-terminal domain (AF-1). Required with NCOA1 to control energy balance between white and brown adipose tissues. Critical regulator of glucose metabolism regulation, acts as RORA coactivator to specifically modulate G6PC1 expression. Involved in the positive regulation of the transcriptional activity of the glucocorticoid receptor NR3C1 by sumoylation enhancer RWDD3. Positively regulates the circadian clock by acting as a transcriptional coactivator for the CLOCK-ARNTL/BMAL1 heterodimer (By similarity).

Cellular Location

Nucleus.

**NCOA2 Blocking Peptide (C-Term) -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)