

**MLL3 Blocking Peptide (N-term)**  
**Synthetic peptide**  
**Catalog # BP22079a****Specification****MLL3 Blocking Peptide (N-term) - Product Information**

Primary Accession [Q8NEZ4](#)  
Other Accession [Q8BRH4](#)

**MLL3 Blocking Peptide (N-term) - Additional Information**

**Gene ID** 58508

**Other Names**

Histone-lysine N-methyltransferase 2C,  
Lysine N-methyltransferase 2C, 2.1.1.43,  
Homologous to ALR protein,  
Myeloid/lymphoid or mixed-lineage  
leukemia protein 3, KMT2C, HALR,  
KIAA1506, MLL3

**Target/Specificity**

The synthetic peptide sequence is selected  
from aa 1162-1175 of HUMAN KMT2C

**Format**

Peptides are lyophilized in a solid powder  
format. Peptides can be reconstituted in  
solution using the appropriate buffer as  
needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6  
months. For long term storage store at  
-20°C.

**Precautions**

This product is for research use only. Not  
for use in diagnostic or therapeutic  
procedures.

**MLL3 Blocking Peptide (N-term) - Protein Information**

**Name** KMT2C

**Synonyms** HALR, KIAA1506, MLL3

**MLL3 Blocking Peptide (N-term) - Background**

Histone methyltransferase. Methylates 'Lys-4'  
of histone H3. H3 'Lys-4' methylation  
represents a specific tag for epigenetic  
transcriptional activation. Central component  
of the MLL2/3 complex, a coactivator complex  
of nuclear receptors, involved in transcriptional  
coactivation. KMT2C/MLL3 may be a catalytic  
subunit of this complex. May be involved in  
leukemogenesis and developmental disorder.

**MLL3 Blocking Peptide (N-term) - References**

Ruault M.,et al.Gene 284:73-81(2002).  
Tan Y.C.,et al.Cancer Detect. Prev.  
25:454-469(2001).  
Hillier L.W.,et al.Nature 424:157-164(2003).  
Nagase T.,et al.DNA Res. 7:143-150(2000).  
Nakajima D.,et al.DNA Res. 9:99-106(2002).

**Function**

Histone methyltransferase that methylates 'Lys-4' of histone H3 (PubMed:<a href="http://www.uniprot.org/citations/22266653" target="\_blank">22266653</a>). H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation. Central component of the MLL2/3 complex, a coactivator complex of nuclear receptors, involved in transcriptional coactivation. KMT2C/MLL3 may be a catalytic subunit of this complex. May be involved in leukemogenesis and developmental disorder.

**Cellular Location**

Nucleus.

**Tissue Location**

Highly expressed in testis and ovary, followed by brain and liver. Also expressed in placenta, peripheral blood, fetal thymus, heart, lung and kidney. Within brain, expression was highest in hippocampus, caudate nucleus, and substantia nigra. Not detected in skeletal muscle and fetal liver

**MLL3 Blocking Peptide (N-term) -  
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)