

Hsp60 Blocking Peptide
Synthetic peptide
Catalog # BP22115a**Specification****Hsp60 Blocking Peptide - Product Information**

Primary Accession [P10809](#)
Other Accession [P31081](#), [Q5NVM5](#)

Hsp60 Blocking Peptide - Additional Information

Gene ID 3329

Other Names

60 kDa heat shock protein, mitochondrial,
60 kDa chaperonin, Chaperonin 60, CPN60,
Heat shock protein 60, HSP-60, Hsp60,
HuCHA60, Mitochondrial matrix protein P1,
P60 lymphocyte protein, HSPD1, HSP60

Target/Specificity

The synthetic peptide sequence is selected
from aa 360-374 of HUMAN HSPD1

Format

Peptides are lyophilized in a solid powder
format. Peptides can be reconstituted in
solution using the appropriate buffer as
needed.

Storage

Maintain refrigerated at 2-8°C for up to 6
months. For long term storage store at
-20°C.

Precautions

This product is for research use only. Not
for use in diagnostic or therapeutic
procedures.

Hsp60 Blocking Peptide - Protein Information

Name HSPD1

Synonyms HSP60

Function

Chaperonin implicated in mitochondrial
protein import and macromolecular

Hsp60 Blocking Peptide - Background

Implicated in mitochondrial protein import
and macromolecular assembly. May facilitate
the correct folding of imported proteins. May
also prevent misfolding and promote the
refolding and proper assembly of unfolded
polypeptides generated under stress
conditions in the mitochondrial matrix.

Hsp60 Blocking Peptide - References

Jindal S.,et al.Mol. Cell. Biol.
9:2279-2283(1989).
Venner T.J.,et al.DNA Cell Biol.
9:545-552(1990).
Hansen J.J.,et al.Hum. Genet. 112:71-77(2003).
Tan J.,et al.Submitted (SEP-2005) to the
EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).

assembly. Together with Hsp10, facilitates the correct folding of imported proteins. May also prevent misfolding and promote the refolding and proper assembly of unfolded polypeptides generated under stress conditions in the mitochondrial matrix (PubMed:1346131, PubMed:11422376). The functional units of these chaperonins consist of heptameric rings of the large subunit Hsp60, which function as a back-to-back double ring. In a cyclic reaction, Hsp60 ring complexes bind one unfolded substrate protein per ring, followed by the binding of ATP and association with 2 heptameric rings of the co-chaperonin Hsp10. This leads to sequestration of the substrate protein in the inner cavity of Hsp60 where, for a certain period of time, it can fold undisturbed by other cell components. Synchronous hydrolysis of ATP in all Hsp60 subunits results in the dissociation of the chaperonin rings and the release of ADP and the folded substrate protein (Probable).

Cellular Location

Mitochondrion matrix.

Hsp60 Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)