

Phospho-ATG13(S355) Blocking Peptide
Synthetic peptide
Catalog # BP3834a**Specification****Phospho-ATG13(S355) Blocking Peptide - Product Information**

Primary Accession [Q75143](#)
Other Accession [Q91YI1](#),
[NP_001136145.1](#),
[NP_055556.2](#)

Phospho-ATG13(S355) Blocking Peptide - Additional Information

Gene ID 9776

Other Names

Autophagy-related protein 13, ATG13,
KIAA0652

Target/Specificity

The synthetic peptide sequence is selected from aa 348-362 of HUMAN ATG13

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-ATG13(S355) Blocking Peptide - Protein Information

Name ATG13

Synonyms KIAA0652

Function

Autophagy factor required for

Phospho-ATG13(S355) Blocking Peptide - Background

Autophagy factor required for autophagosome formation. Target of the TOR kinase signaling pathway that regulates autophagy through the control of the phosphorylation status of ATG13 and ULK1, and the regulation of the ATG13-ULK1-RB1CC1 complex.

Phospho-ATG13(S355) Blocking Peptide - References

Ferreira, R.C., et al. Nat. Genet. 42(9):777-780(2010)
Hosokawa, N., et al. Autophagy 5(7):973-979(2009)
Mercer, C.A., et al. Autophagy 5(5):649-662(2009)
Ganley, I.G., et al. J. Biol. Chem. 284(18):12297-12305(2009)
Chan, E.Y., et al. Mol. Cell. Biol. 29(1):157-171(2009)

autophagosome formation and mitophagy. Target of the TOR kinase signaling pathway that regulates autophagy through the control of the phosphorylation status of ATG13 and ULK1, and the regulation of the ATG13-ULK1-RB1CC1 complex. Through its regulation of ULK1 activity, plays a role in the regulation of the kinase activity of mTORC1 and cell proliferation.

Cellular Location

Cytoplasm, cytosol. Preautophagosomal structure. Note=Under starvation conditions, is localized to punctate structures primarily representing the isolation membrane; the isolation membrane sequesters a portion of the cytoplasm resulting in autophagosome formation

Phospho-ATG13(S355) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)