

**Phospho-mouse BAD(S96) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP3849a****Specification****Phospho-mouse BAD(S96) Blocking Peptide - Product Information**

Primary Accession [Q61337](#)  
Other Accession [NP\\_031548.1](#)

**Phospho-mouse BAD(S96) Blocking Peptide - Additional Information**

**Gene ID** 12015

**Other Names**

Bcl2-associated agonist of cell death, BAD,  
Bcl-2-binding component 6,  
Bcl-xL/Bcl-2-associated death promoter,  
Bcl2 antagonist of cell death, Bad, Bbc6

**Target/Specificity**

The synthetic peptide sequence is selected from aa 90-102 of MOUSE Bad

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Phospho-mouse BAD(S96) Blocking Peptide - Protein Information**

**Name** Bad

**Synonyms** Bbc6

**Function**

Promotes cell death. Successfully competes

**Phospho-mouse BAD(S96) Blocking Peptide - Background**

BAD promotes cell death. Successfully competes for the binding to Bcl-X(L), Bcl-2 and Bcl-W, thereby affecting the level of heterodimerization of these proteins with BAX. Can reverse the death repressor activity of Bcl-X(L), but not that of Bcl-2. Appears to act as a link between growth factor receptor signaling and the apoptotic pathways.

**Phospho-mouse BAD(S96) Blocking Peptide - References**

Santidrian, A.F., et al. Blood 116(16):3023-3032(2010)  
Frenzel, A., et al. Blood 115(5):995-1005(2010)  
Quoyer, J., et al. J. Biol. Chem. 285(3):1989-2002(2010)  
Polzien, L., et al. J. Biol. Chem. 284(41):28004-28020(2009)  
Wu, X., et al. Diabetologia 52(10):2130-2141(2009)

for the binding to Bcl-X(L), Bcl-2 and Bcl-W, thereby affecting the level of heterodimerization of these proteins with BAX. Can reverse the death repressor activity of Bcl-X(L), but not that of Bcl-2. Appears to act as a link between growth factor receptor signaling and the apoptotic pathways.

**Cellular Location**

Mitochondrion outer membrane. Cytoplasm.

Note=Colocalizes with HIF3A isoform 2 in the cytoplasm (PubMed:21546903). Upon phosphorylation, locates to the cytoplasm.

**Phospho-mouse BAD(S96) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)