

**Phospho-mouse ERBB2(S1114) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP3850a****Specification****Phospho-mouse ERBB2(S1114) Blocking Peptide - Product Information**

Primary Accession [P70424](#)  
Other Accession [P06494](#),  
[NP\\_001003817.1](#)

**Phospho-mouse ERBB2(S1114) Blocking Peptide - Additional Information**

**Gene ID** 13866

**Other Names**

Receptor tyrosine-protein kinase erbB-2,  
Proto-oncogene Neu, Proto-oncogene  
c-ErbB-2, p185erbB2, CD340, Erbb2,  
Kiaa3023, Neu

**Target/Specificity**

The synthetic peptide sequence is selected  
from aa 1107-1120 of MOUSE Erbb2

**Format**

Peptides are lyophilized in a solid powder  
format. Peptides can be reconstituted in  
solution using the appropriate buffer as  
needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6  
months. For long term storage store at  
-20°C.

**Precautions**

This product is for research use only. Not  
for use in diagnostic or therapeutic  
procedures.

**Phospho-mouse ERBB2(S1114) Blocking Peptide - Protein Information**

**Name** Erbb2

**Synonyms** Kiaa3023, Neu

**Function**

**Phospho-mouse ERBB2(S1114) Blocking Peptide - Background**

This gene encodes a member of the epidermal  
growth factor  
(EGF) receptor family of receptor tyrosine  
kinases. This protein  
has no ligand binding domain of its own and  
therefore cannot bind  
growth factors. However, it does bind tightly to  
other ligand-bound  
EGF receptor family members to form a  
heterodimer, stabilizing  
ligand binding and enhancing kinase-mediated  
activation of  
downstream signalling pathways, such as  
those involving  
mitogen-activated protein kinase and  
phosphatidylinositol-3 kinase.  
Allelic variations at amino acid positions 654  
and 655 of isoform a  
(positions 624 and 625 of isoform b) have been  
reported, with the  
most common allele, Ile654/Ile655, shown  
here. Amplification and/or  
overexpression of this gene has been reported  
in numerous cancers,  
including breast and ovarian tumors.  
Alternative splicing results  
in several additional transcript variants, some  
encoding different  
isoforms and others that have not been fully  
characterized.

**Phospho-mouse ERBB2(S1114) Blocking Peptide - References**

Cabodi, S., et al. FASEB J.  
24(10):3796-3808(2010)  
Johnson, E., et al. J. Biol. Chem.  
285(38):29491-29501(2010)  
Huck, L., et al. Proc. Natl. Acad. Sci. U.S.A.  
107(35):15559-15564(2010)  
Chuang, T.D., et al. J. Biol. Chem.  
285(31):23598-23606(2010)  
Simeone, L., et al. J. Neurosci.  
30(19):6620-6634(2010)

Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization (By similarity).

#### **Cellular Location**

Cell membrane

{ECO:0000250|UniProtKB:P04626};

Single-pass type I membrane protein

{ECO:0000250|UniProtKB:P04626} Early endosome

{ECO:0000250|UniProtKB:P04626}.

Cytoplasm, perinuclear region

{ECO:0000250|UniProtKB:P04626}. Nucleus

{ECO:0000250|UniProtKB:P04626}.

Note=Translocation to the nucleus requires endocytosis, probably endosomal sorting and is mediated by importin beta-1/KPNB1.

Also detected in endosome-to-TGN retrograde vesicles.

{ECO:0000250|UniProtKB:P04626}

#### **Tissue Location**

Expressed predominantly in uterine epithelial cells. In the muscle, expression localizes to the synaptic sites of muscle fibers

#### **Phospho-mouse ERBB2(S1114) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)