

Phospho-TSC1(T310) Blocking Peptide
Synthetic peptide
Catalog # BP3865a**Specification****Phospho-TSC1(T310) Blocking Peptide - Product Information**

Primary Accession [Q92574](#)
Other Accession [NP_000359.1](#)

Phospho-TSC1(T310) Blocking Peptide - Additional Information

Gene ID 7248

Other Names

Hamartin, Tuberous sclerosis 1 protein, TSC1, KIAA0243, TSC

Target/Specificity

The synthetic peptide sequence is selected from aa 304-316 of HUMAN TSC1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-TSC1(T310) Blocking Peptide - Protein Information

Name TSC1

Synonyms KIAA0243, TSC

Function

In complex with TSC2, inhibits the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1

Phospho-TSC1(T310) Blocking Peptide - Background

This gene encodes a growth inhibitory protein thought to play a role in the stabilization of tuberin. Mutations in this gene have been associated with tuberous sclerosis. Alternative splicing results in multiple transcript variants.

Phospho-TSC1(T310) Blocking Peptide - References

Hoogeveen-Westerveld, M., et al. Biochim. Biophys. Acta 1802(9):774-781(2010)
Mehta, M.S., et al. Breast Cancer Res. Treat. (2010) In press :
Mieulet, V., et al. Trends Mol Med 16(7):329-335(2010)
Liu, C.Y., et al. Carcinogenesis 31(7):1259-1263(2010)
Guo, L., et al. Acta Biochim. Biophys. Sin. (Shanghai) 42(4):266-273(2010)

and EIF4EBP1 by negatively regulating mTORC1 signaling (PubMed:12271141, PubMed:28215400). Seems not to be required for TSC2 GAP activity towards RHEB (PubMed:15340059). Implicated as a tumor suppressor. Involved in microtubule-mediated protein transport, but this seems to be due to unregulated mTOR signaling (By similarity). Acts as a co-chaperone for HSP90AA1 facilitating HSP90AA1 chaperoning of protein clients such as kinases, TSC2 and glucocorticoid receptor NR3C1 (PubMed:29127155). Increases ATP binding to HSP90AA1 and inhibits HSP90AA1 ATPase activity (PubMed:29127155). Competes with the activating co-chaperone AHSA1 for binding to HSP90AA1, thereby providing a reciprocal regulatory mechanism for chaperoning of client proteins (PubMed:29127155). Recruits TSC2 to HSP90AA1 and stabilizes TSC2 by preventing the interaction between TSC2 and ubiquitin ligase HERC1 (PubMed:16464865, PubMed:29127155).

Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein. Note=At steady state found in association with membranes.

Tissue Location

Highly expressed in skeletal muscle, followed by heart, brain, placenta, pancreas, lung, liver and kidney. Also expressed in embryonic kidney cells

Phospho-TSC1(T310) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)