

**Phospho-mouse p21Cip1(S125) Blocking Peptide**

Synthetic peptide

Catalog # BP3875a

**Specification****Phospho-mouse p21Cip1(S125) Blocking Peptide  
- Product Information**

Primary Accession [P39689](#)  
Other Accession [NP\\_001129489.1](#)

**Phospho-mouse p21Cip1(S125) Blocking Peptide  
- Additional Information****Gene ID** 12575**Other Names**

Cyclin-dependent kinase inhibitor 1,  
CDK-interacting protein 1, Melanoma  
differentiation-associated protein, p21,  
Cdkn1a, Cip1, Waf1

**Target/Specificity**

The synthetic peptide sequence is selected  
from aa 119-132 of MOUSE Cdkn1a

**Format**

Peptides are lyophilized in a solid powder  
format. Peptides can be reconstituted in  
solution using the appropriate buffer as  
needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6  
months. For long term storage store at  
-20°C.

**Precautions**

This product is for research use only. Not  
for use in diagnostic or therapeutic  
procedures.

**Phospho-mouse p21Cip1(S125) Blocking Peptide  
- Protein Information****Name** Cdkn1a**Synonyms** Cip1, Waf1**Function**

May be involved in p53/TP53 mediated

**Phospho-mouse p21Cip1(S125) Blocking  
Peptide - Background**

The protein encoded by this gene belongs to  
the highly  
conserved cyclin family, whose members are  
characterized by a  
dramatic periodicity in protein abundance  
through the cell cycle.  
Cyclins function as regulators of CDK kinases.  
Different cyclins  
exhibit distinct expression and degradation  
patterns which  
contribute to the temporal coordination of each  
mitotic event. This  
cyclin forms a complex with and functions as a  
regulatory subunit  
of CDK4 or CDK6, whose activity is required for  
cell cycle G1/S  
transition. This protein has been shown to  
interact with and be  
involved in the phosphorylation of tumor  
suppressor protein Rb. The  
CDK4 activity associated with this cyclin was  
reported to be  
necessary for cell cycle progression through G2  
phase into mitosis  
after UV radiation. Several transcript variants  
encoding different  
isoforms have been found for this gene.

**Phospho-mouse p21Cip1(S125) Blocking  
Peptide - References**

Liu, C.Y., et al. Carcinogenesis  
31(7):1259-1263(2010)  
Kim, J., et al. Cytokine 50(1):42-49(2010)  
Kamatani, Y., et al. Nat. Genet.  
42(3):210-215(2010)  
Gumina, M.R., et al. Cell Cycle  
9(4):820-828(2010)  
Radulovich, N., et al. Mol. Cancer 9, 24 (2010) :

inhibition of cellular proliferation in response to DNA damage. Binds to and inhibits cyclin- dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D-CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D- CDK4 complex (PubMed:<a href="http://www.uniprot.org/citations/25329316" target="\_blank">25329316</a>). Inhibits DNA synthesis by DNA polymerase delta by competing with POLD3 for PCNA binding (By similarity). Plays an important role in controlling cell cycle progression and DNA damage-induced G2 arrest (By similarity).

**Cellular Location**

Cytoplasm

{ECO:0000250|UniProtKB:P38936}. Nucleus

**Phospho-mouse p21Cip1(S125) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)