

Phospho-p38 MAPK (Y182) Blocking Peptide
Synthetic peptide
Catalog # BP3905a**Specification****Phospho-p38 MAPK (Y182) Blocking Peptide -
Product Information**

Primary Accession [Q15759](#)
Other Accession [Q9WUI1](#)

**Phospho-p38 MAPK (Y182) Blocking Peptide -
Additional Information**

Gene ID 5600

Other Names

Mitogen-activated protein kinase 11, MAP
kinase 11, MAPK 11, 2.7.11.24,
Mitogen-activated protein kinase p38 beta,
MAP kinase p38 beta, p38b,
Stress-activated protein kinase 2b, SAPK2b,
p38-2, MAPK11, PRKM11, SAPK2, SAPK2B

Target/Specificity

The synthetic peptide sequence is selected
from aa 175-186 of HUMAN MAPK11

Format

Peptides are lyophilized in a solid powder
format. Peptides can be reconstituted in
solution using the appropriate buffer as
needed.

Storage

Maintain refrigerated at 2-8°C for up to 6
months. For long term storage store at
-20°C.

Precautions

This product is for research use only. Not
for use in diagnostic or therapeutic
procedures.

**Phospho-p38 MAPK (Y182) Blocking Peptide -
Protein Information**

Name MAPK11

Synonyms PRKM11, SAPK2, SAPK2B

**Phospho-p38 MAPK (Y182) Blocking
Peptide - Background**

Serine/threonine kinase which acts as an
essential component of the MAP kinase signal
transduction pathway. MAPK11 is one of the
four p38 MAPKs which play an important role in
the cascades of cellular responses evoked by
extracellular stimuli such as proinflammatory
cytokines or physical stress leading to direct
activation of transcription factors. Accordingly,
p38 MAPKs phosphorylate a broad range of
proteins and it has been estimated that they
may have approximately 200 to 300 substrates
each. MAPK11 functions are mostly redundant
with those of MAPK14. Some of the targets are
downstream kinases which are activated
through phosphorylation and further
phosphorylate additional targets.
RPS6KA5/MSK1 and RPS6KA4/MSK2 can
directly phosphorylate and activate
transcription factors such as CREB1, ATF1, the
NF-kappa-B isoform RELA/NFKB3, STAT1 and
STAT3, but can also phosphorylate histone H3
and the nucleosomal protein HMGN1.
RPS6KA5/MSK1 and RPS6KA4/MSK2 play
important roles in the rapid induction of
immediate-early genes in response to stress or
mitogenic stimuli, either by inducing chromatin
remodeling or by recruiting the transcription
machinery. On the other hand, two other
kinase targets, MAPKAPK2/MK2 and
MAPKAPK3/MK3, participate in the control of
gene expression mostly at the
post-transcriptional level, by phosphorylating
ZFP36 (tristetraprolin) and ELAVL1, and by
regulating EEF2K, which is important for the
elongation of mRNA during translation.
MKNK1/MNK1 and MKNK2/MNK2, two other
kinases activated by p38 MAPKs, regulate
protein synthesis by phosphorylating the
initiation factor EIF4E2. In the cytoplasm, the
p38 MAPK pathway is an important regulator of
protein turnover. For example, CFLAR is an
inhibitor of TNF-induced apoptosis whose
proteasome-mediated degradation is regulated
by p38 MAPK phosphorylation. Ectodomain
shedding of transmembrane proteins is

Function

Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK11 is one of the four p38 MAPKs which play an important role in the cascades of cellular responses evoked by extracellular stimuli such as proinflammatory cytokines or physical stress leading to direct activation of transcription factors. Accordingly, p38 MAPKs phosphorylate a broad range of proteins and it has been estimated that they may have approximately 200 to 300 substrates each. MAPK11 functions are mostly redundant with those of MAPK14. Some of the targets are downstream kinases which are activated through phosphorylation and further phosphorylate additional targets. RPS6KA5/MSK1 and RPS6KA4/MSK2 can directly phosphorylate and activate transcription factors such as CREB1, ATF1, the NF-kappa-B isoform RELA/NFKB3, STAT1 and STAT3, but can also phosphorylate histone H3 and the nucleosomal protein HMGN1. RPS6KA5/MSK1 and RPS6KA4/MSK2 play important roles in the rapid induction of immediate-early genes in response to stress or mitogenic stimuli, either by inducing chromatin remodeling or by recruiting the transcription machinery. On the other hand, two other kinase targets, MAPKAPK2/MK2 and MAPKAPK3/MK3, participate in the control of gene expression mostly at the post-transcriptional level, by phosphorylating ZFP36 (tristetraprolin) and ELAVL1, and by regulating EEF2K, which is important for the elongation of mRNA during translation. MKNK1/MNK1 and MKNK2/MNK2, two other kinases activated by p38 MAPKs, regulate protein synthesis by phosphorylating the initiation factor EIF4E2. In the cytoplasm, the p38 MAPK pathway is an important regulator of protein turnover. For example, CFLAR is an inhibitor of TNF-induced apoptosis whose proteasome-mediated degradation is regulated by p38 MAPK phosphorylation. Ectodomain shedding of transmembrane proteins is regulated by p38 MAPKs as well. In response to inflammatory stimuli, p38 MAPKs phosphorylate the membrane-associated metalloprotease ADAM17. Such phosphorylation is required for ADAM17-mediated ectodomain shedding of TGF-alpha family ligands, which results in the activation of EGFR signaling and cell

regulated by p38 MAPKs as well. In response to inflammatory stimuli, p38 MAPKs phosphorylate the membrane-associated metalloprotease ADAM17. Such phosphorylation is required for ADAM17-mediated ectodomain shedding of TGF-alpha family ligands, which results in the activation of EGFR signaling and cell proliferation. Additional examples of p38 MAPK substrates are the FGFR1. FGFR1 can be translocated from the extracellular space into the cytosol and nucleus of target cells, and regulates processes such as rRNA synthesis and cell growth. FGFR1 translocation requires p38 MAPK activation. In the nucleus, many transcription factors are phosphorylated and activated by p38 MAPKs in response to different stimuli. Classical examples include ATF1, ATF2, ATF6, ELK1, PTPRH, DDIT3, TP53/p53 and MEF2C and MEF2A. The p38 MAPKs are emerging as important modulators of gene expression by regulating chromatin modifiers and remodelers. The promoters of several genes involved in the inflammatory response, such as IL6, IL8 and IL12B, display a p38 MAPK-dependent enrichment of histone H3 phosphorylation on 'Ser-10' (H3S10ph) in LPS-stimulated myeloid cells. This phosphorylation enhances the accessibility of the cryptic NF-kappa-B-binding sites marking promoters for increased NF-kappa-B recruitment.

Phospho-p38 MAPK (Y182) Blocking Peptide - References

- Jiang Y., et al. J. Biol. Chem. 271:17920-17926(1996).
Jiang Y., et al. Submitted (APR-1997) to the EMBL/GenBank/DDBJ databases.
Kumar S., et al. Biochem. Biophys. Res. Commun. 235:533-538(1997).
Enslen H., et al. J. Biol. Chem. 273:1741-1748(1998).
Goedert M., et al. EMBO J. 16:3563-3571(1997).

proliferation. Additional examples of p38 MAPK substrates are the FGFR1. FGFR1 can be translocated from the extracellular space into the cytosol and nucleus of target cells, and regulates processes such as rRNA synthesis and cell growth. FGFR1 translocation requires p38 MAPK activation. In the nucleus, many transcription factors are phosphorylated and activated by p38 MAPKs in response to different stimuli. Classical examples include ATF1, ATF2, ATF6, ELK1, PTPRH, DDIT3, TP53/p53 and MEF2C and MEF2A. The p38 MAPKs are emerging as important modulators of gene expression by regulating chromatin modifiers and remodelers. The promoters of several genes involved in the inflammatory response, such as IL6, IL8 and IL12B, display a p38 MAPK-dependent enrichment of histone H3 phosphorylation on 'Ser-10' (H3S10ph) in LPS-stimulated myeloid cells. This phosphorylation enhances the accessibility of the cryptic NF-kappa-B-binding sites marking promoters for increased NF-kappa-B recruitment.

Cellular Location

Cytoplasm. Nucleus.

Tissue Location

Highest levels in the brain and heart. Also expressed in the placenta, lung, liver, skeletal muscle, kidney and pancreas

Phospho-p38 MAPK (Y182) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)