

CA9 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP5000a**Specification****CA9 Antibody (N-term) Blocking Peptide - Product Information**Primary Accession [Q16790](#)**CA9 Antibody (N-term) Blocking Peptide - Additional Information**

Gene ID 768

Other Names

Carbonic anhydrase 9, Carbonate dehydratase IX, Carbonic anhydrase IX, CA-IX, CAIX, Membrane antigen MN, P54/58N, Renal cell carcinoma-associated antigen G250, RCC-associated antigen G250, pMW1, CA9, G250, MN

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP5000a](/product/products/AP5000a) was selected from the N-term region of human CA9. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CA9 Antibody (N-term) Blocking Peptide - Protein Information**CA9 Antibody (N-term) Blocking Peptide - Background**

Carbonic anhydrases (CAs) are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, and the formation of aqueous humor, cerebrospinal fluid, saliva, and gastric acid. They show extensive diversity in tissue distribution and in their subcellular localization. CA IX is a transmembrane protein and the only tumor-associated carbonic anhydrase isoenzyme known. It is expressed in all clear-cell renal cell carcinoma, but is not detected in normal kidney or most other normal tissues. It may be involved in cell proliferation and transformation. This gene was mapped to 17q21.2 by fluorescence in situ hybridization, however, radiation hybrid mapping localized it to 9p13-p12.

CA9 Antibody (N-term) Blocking Peptide - References

Grabmaier, K., et al., *Oncogene* 23(33):5624-5631 (2004). Kaluzova, M., et al., *Mol. Cell. Biol.* 24(13):5757-5766 (2004). Span, P.N., et al., *Br. J. Cancer* 89(2):271-276 (2003). Hedley, D., et al., *Clin. Cancer Res.* 9(15):5666-5674 (2003). Bui, M.H., et al., *Clin. Cancer Res.* 9(2):802-811 (2003).

Name CA9

Synonyms G250, MN

Function

Reversible hydration of carbon dioxide. Participates in pH regulation. May be involved in the control of cell proliferation and transformation. Appears to be a novel specific biomarker for a cervical neoplasia.

Cellular Location

Nucleus. Nucleus, nucleolus. Cell membrane; Single-pass type I membrane protein. Cell projection, microvillus membrane; Single-pass type I membrane protein. Note=Found on the surface microvilli and in the nucleus, particularly in nucleolus

Tissue Location

Expressed primarily in carcinoma cells lines. Expression is restricted to very few normal tissues and the most abundant expression is found in the epithelial cells of gastric mucosa

CA9 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)