

FE65 (APBB1) Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP6100a**Specification****FE65 (APBB1) Antibody (C-term) Blocking peptide**
- Product Information

Primary Accession [O00213](#)
Other Accession [NP_663722](#)

FE65 (APBB1) Antibody (C-term) Blocking peptide
- Additional Information

Gene ID 322

Other Names

Amyloid beta A4 precursor protein-binding family B member 1, Protein Fe65, APBB1, FE65, RIR

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6100a](/product/products/AP6100a) was selected from the C-term region of human APBB1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

FE65 (APBB1) Antibody (C-term) Blocking peptide
- Protein Information

Name APBB1

FE65 (APBB1) Antibody (C-term) Blocking peptide - Background

APBB1 is a member of the Fe65 protein family. It is an adaptor protein localized in the nucleus. It interacts with the Alzheimer's disease amyloid precursor protein (APP), transcription factor CP2/LSF/LBP1 and the low-density lipoprotein receptor-related protein. APP functions as a cytosolic anchoring site that can prevent the gene product's nuclear translocation. This encoded protein could play an important role in the pathogenesis of Alzheimer's disease. It is thought to regulate transcription. Also it is observed to block cell cycle progression by downregulating thymidylate synthase expression.

FE65 (APBB1) Antibody (C-term) Blocking peptide - References

Kinoshita, A., et al., J. Biol. Chem. 278(42):41182-41188 (2003). Walsh, D.M., et al., Biochemistry 42(22):6664-6673 (2003). Zhao, Q., et al., Biochemistry 42(12):3627-3634 (2003). Hu, Q., et al., Hum. Mol. Genet. 11(4):465-475 (2002). Bruni, P., et al., J. Biol. Chem. 277(38):35481-35488 (2002).

Synonyms FE65, RIR**Function**

Transcription coregulator that can have both coactivator and corepressor functions. Adapter protein that forms a transcriptionally active complex with the gamma-secretase-derived amyloid precursor protein (APP) intracellular domain. Plays a central role in the response to DNA damage by translocating to the nucleus and inducing apoptosis. May act by specifically recognizing and binding histone H2AX phosphorylated on 'Tyr-142' (H2AXY142ph) at double-strand breaks (DSBs), recruiting other pro-apoptosis factors such as MAPK8/JNK1. Required for histone H4 acetylation at double-strand breaks (DSBs). Its ability to specifically bind modified histones and chromatin modifying enzymes such as KAT5/TIP60, probably explains its transcription activation activity. Functions in association with TSHZ3, SET and HDAC factors as a transcriptional repressor, that inhibits the expression of CASP4. Associates with chromatin in a region surrounding the CASP4 transcriptional start site(s).

Cellular Location

Cell membrane. Cytoplasm. Nucleus. Cell projection, growth cone. Nucleus speckle
Note=Colocalizes with TSHZ3 in axonal growth cone (By similarity). In normal conditions, it mainly localizes to the cytoplasm, while a small fraction is tethered to the cell membrane via its interaction with APP. Following exposure to DNA damaging agents, it is released from cell membrane and translocates to the nucleus. Nuclear translocation is under the regulation of APP. Colocalizes with TSHZ3 in the nucleus. Colocalizes with NEK6 at the nuclear speckles. Phosphorylation at Ser- 610 by SGK1 promotes its localization to the nucleus (By similarity)

Tissue Location

Highly expressed in brain; strongly reduced in post-mortem elderly subjects with Alzheimer disease

FE65 (APBB1) Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you

may find useful for product applications.

- [Blocking Peptides](#)