

Parp12 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP6298b**Specification****Parp12 Antibody (C-term) Blocking Peptide - Product Information**

Primary Accession [Q8BZ20](#)
Other Accession [NP_766481](#)

Parp12 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 243771

Other Names

Poly [ADP-ribose] polymerase 12, PARP-12, ADP-ribosyltransferase diphtheria toxin-like 12, ARTD12, Zinc finger CCCH domain-containing protein 1, Parp12, Zc3hdc1

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6298b](#) was selected from the C-term region of human Parp12 . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Parp12 Antibody (C-term) Blocking Peptide - Protein Information**Parp12 Antibody (C-term) Blocking Peptide - Background**

Poly(ADP-ribosyl)ation is an immediate DNA-damage-dependent post-translational modification of histones and other nuclear proteins that contributes to the survival of injured proliferating cells. Poly(ADP-ribose) polymerases (PARPs) now constitute a large family of 18 proteins, encoded by different genes and displaying a conserved catalytic domain in which PARP-1 (113 kDa), the founding member, and PARP-2 (62 kDa) are so far the sole enzymes whose catalytic activity has been shown to be immediately stimulated by DNA strand breaks. A large repertoire of sequences encoding novel PARPs now extends considerably the field of poly(ADP-ribosyl)ation reactions to various aspects of the cell biology including cell proliferation and cell death. Some of these new members interact with each other, share common partners and common subcellular localizations suggesting possible fine tuning in the regulation of this post-translational modification of proteins.

Parp12 Antibody (C-term) Blocking Peptide - References

Bailey,P.J., Exp. Cell Res. 312 (16), 3108-3119 (2006)Kato,M., Int. J. Oncol. 23 (2), 541-547 (2003)

Name Parp12
{ECO:0000312|MGI:MGI:2143990}

Synonyms Zc3hdc1

Function
Mono-ADP-ribosyltransferase that mediates
mono-ADP- ribosylation of target proteins.

Cellular Location
Nucleus.

Parp12 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Blocking Peptides](#)