

**NDUFS4 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP6932b****Specification****NDUFS4 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [O43181](#)**NDUFS4 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 4724**Other Names**

NADH dehydrogenase [ubiquinone] iron-sulfur protein 4, mitochondrial, Complex I-18 kDa, CI-18 kDa, Complex I-AQDQ, CI-AQDQ, NADH-ubiquinone oxidoreductase 18 kDa subunit, NDUFS4

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP6932b](/products/AP6932b) was selected from the C-term region of human NDUFS4. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**NDUFS4 Antibody (C-term) Blocking Peptide - Protein Information****NDUFS4 Antibody (C-term) Blocking Peptide - Background**

NDUFS4 is an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase(Complex I), or NADH:ubiquinone oxidoreductase, the first multi-subunit enzyme complex of the mitochondrial respiratory chain. Complex I plays a vital role in cellular ATP production, the primary source of energy for many crucial processes in living cells. It removes electrons from NADH and passes them by a series of different protein-coupled redox centers to the electron acceptor ubiquinone. In well-coupled mitochondria, the electron flux leads to ATP generation via the building of a proton gradient across the inner membrane.

**NDUFS4 Antibody (C-term) Blocking Peptide - References**

Panelli,D., et.al., Biochimie 90 (10), 1452-1460 (2008)

**Name** NDUFS4

**Function**

Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

**Cellular Location**

Mitochondrion inner membrane; Peripheral membrane protein; Matrix side. Note=The interaction with BCAP31 mediates mitochondria localization.

**NDUFS4 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)