

**AD\_K2 Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP7005a****Specification****AD\_K2 Antibody (C-term) Blocking peptide -  
Product Information**Primary Accession [P35626](#)**AD\_K2 Antibody (C-term) Blocking peptide -  
Additional Information****Gene ID 157****Other Names**Beta-adrenergic receptor kinase 2,  
Beta-ARK-2, G-protein-coupled receptor  
kinase 3, ADRBK2, BARK2, GRK3**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP7005a](/product/products/AP7005a) was selected from the C-term region of human GRK3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**AD\_K2 Antibody (C-term) Blocking peptide -  
Protein Information****Name** GRK3 {ECO:0000312|MIM:109636}**AD\_K2 Antibody (C-term) Blocking peptide  
- Background**

The beta-adrenergic receptor kinase specifically phosphorylates the agonist-occupied form of the beta-adrenergic and related G protein-coupled receptors. Overall, the ADRBK2 enzyme, also known as GRK3, has 85% amino acid similarity with ADRBK1, with the protein kinase catalytic domain having 95% similarity. The ADRBK2 mRNA is approximately 8 kilobases with a distribution similar to that of ADRBK1. These data suggest the existence of a family of receptor kinases which may serve broadly to regulate receptor function.

**AD\_K2 Antibody (C-term) Blocking peptide  
- References**

Calabrese, G., et al., Genomics 23(1):286-288 (1994). Parruti, G., et al., Biochem. Biophys. Res. Commun. 190(2):475-481 (1993). Benovic, J.L., et al., J. Biol. Chem. 266(23):14939-14946 (1991).

**Synonyms** ADRBK2, BARK2

**Function**

Specifically phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors.

**AD\_K2 Antibody (C-term) Blocking peptide  
- Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)