

PDK1 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP7038a**Specification****PDK1 Antibody (N-term) Blocking Peptide -
Product Information**Primary Accession [Q15118](#)**PDK1 Antibody (N-term) Blocking Peptide -
Additional Information****Gene ID** 5163**Other Names**

[Pyruvate dehydrogenase
(acetyl-transferring)] kinase isozyme 1,
mitochondrial, Pyruvate dehydrogenase
kinase isoform 1, PDH kinase 1, PDK1,
PDHK1

Target/Specificity

The synthetic peptide sequence used to
generate the antibody [AP7038a](/product/products/AP7038a) was
selected from the N-term region of human
PDK1 . A 10 to 100 fold molar excess to
antibody is recommended. Precise
conditions should be optimized for a
particular assay.

Format

Peptides are lyophilized in a solid powder
format. Peptides can be reconstituted in
solution using the appropriate buffer as
needed.

Storage

Maintain refrigerated at 2-8°C for up to 6
months. For long term storage store at
-20°C.

Precautions

This product is for research use only. Not
for use in diagnostic or therapeutic
procedures.

**PDK1 Antibody (N-term) Blocking Peptide -
Protein Information****PDK1 Antibody (N-term) Blocking Peptide
- Background**

Pyruvate dehydrogenase (PDH) is a
mitochondrial multienzyme complex that
catalyzes the oxidative decarboxylation of
pyruvate and is one of the major enzymes
responsible for the regulation of homeostasis
of carbohydrate fuels in mammals. The
enzymatic activity is regulated by a
phosphorylation/dephosphorylation cycle.
Phosphorylation of PDH by a specific pyruvate
dehydrogenase kinase (PDK) results in
inactivation.

**PDK1 Antibody (N-term) Blocking Peptide
- References**

Sato, S., et al., J. Biol. Chem.
277(42):39360-39367 (2002).Frodin, M., et al.,
EMBO J. 21(20):5396-5407 (2002).King, C.C., et
al., J. Biol. Chem. 275(24):18108-18113
(2000).Gudi, R., et al., J. Biol. Chem.
270(48):28989-28994 (1995).

Name PDK1

Synonyms PDHK1

Function

Kinase that plays a key role in regulation of glucose and fatty acid metabolism and homeostasis via phosphorylation of the pyruvate dehydrogenase subunits PDHA1 and PDHA2. This inhibits pyruvate dehydrogenase activity, and thereby regulates metabolite flux through the tricarboxylic acid cycle, down-regulates aerobic respiration and inhibits the formation of acetyl-coenzyme A from pyruvate. Plays an important role in cellular responses to hypoxia and is important for cell proliferation under hypoxia. Protects cells against apoptosis in response to hypoxia and oxidative stress.

Cellular Location

Mitochondrion matrix

Tissue Location

Expressed predominantly in the heart.
Detected at lower levels in liver, skeletal muscle and pancreas

**PDK1 Antibody (N-term) Blocking Peptide
- Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)