

SMYD3 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP7344a**Specification****SMYD3 Antibody (N-term) Blocking Peptide -
Product Information**Primary Accession [Q9H7B4](#)**SMYD3 Antibody (N-term) Blocking Peptide -
Additional Information****Gene ID** 64754**Other Names**Histone-lysine N-methyltransferase SMYD3,
SET and MYND domain-containing protein 3,
Zinc finger MYND domain-containing protein
1, SMYD3, ZMYND1, ZNFN3A1**Target/Specificity**

The synthetic peptide sequence used to
generate the antibody [AP7344a](/products/AP7344a)
was selected from the N-term region of
human SMYD3. A 10 to 100 fold molar
excess to antibody is recommended.
Precise conditions should be optimized for a
particular assay.

FormatPeptides are lyophilized in a solid powder
format. Peptides can be reconstituted in
solution using the appropriate buffer as
needed.**Storage**Maintain refrigerated at 2-8°C for up to 6
months. For long term storage store at
-20°C.**Precautions**This product is for research use only. Not
for use in diagnostic or therapeutic
procedures.**SMYD3 Antibody (N-term) Blocking Peptide -
Protein Information****Name** SMYD3**SMYD3 Antibody (N-term) Blocking
Peptide - Background**SMYD3 is a histone methyltransferase that
plays a role in transcriptional regulation as a
member of an RNA polymerase complex.**SMYD3 Antibody (N-term) Blocking
Peptide - References**

Zou,J.N., Wang,S.Z. Cancer Lett. 280 (1),
78-85 (2009)Wang,S.Z., Luo,X.G. BMB Rep 41
(4), 294-299 (2008)Wang,H., Liu,Y. Cancer Sci.
99 (4), 787-791 (2008)Barlesi,F., Giaccone,G.
Int. J. Cancer 122 (6), 1441-1442
(2008)Hamamoto,R., Silva,F.P. Cancer Sci. 97
(2), 113-118 (2006)

Synonyms ZMYND1, ZNFN3A1**Function**

Histone methyltransferase. Specifically methylates 'Lys-4' of histone H3, inducing di- and tri-methylation, but not monomethylation (PubMed:15235609, PubMed:22419068). Also methylates 'Lys-5' of histone H4 (PubMed:22419068). Plays an important role in transcriptional activation as a member of an RNA polymerase complex (PubMed:15235609). Binds DNA containing 5'-CCCTCC-3' or 5'-GAGGGG-3' sequences (PubMed:15235609).

Cellular Location

Cytoplasm. Nucleus. Note=Mainly cytoplasmic when cells are arrested at G0/G1. Accumulates in the nucleus at S phase and G2/M.

Tissue Location

Expressed in skeletal muscles and testis. Overexpressed in a majority of colorectal and hepatocellular carcinomas.

SMYD3 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)