

**B-RAF Antibody (S445) Blocking peptide**  
Synthetic peptide  
Catalog # BP7810f**Specification****B-RAF Antibody (S445) Blocking peptide -  
Product Information**Primary Accession [P15056](#)**B-RAF Antibody (S445) Blocking peptide -  
Additional Information**

Gene ID 673

**Other Names**Serine/threonine-protein kinase B-raf,  
Proto-oncogene B-Raf, p94, v-Raf murine  
sarcoma viral oncogene homolog B1, BRAF,  
BRAF1, RAFB1**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP7810f](#) was selected from the S445 region of human BRAF. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**B-RAF Antibody (S445) Blocking peptide - Protein  
Information**Name BRAF ([HGNC:1097](#))**B-RAF Antibody (S445) Blocking peptide -  
Background**

BRAF, a member of the RAF subfamily of Ser/Thr protein kinases, is involved in the transduction of mitogenic signals from the cell membrane to the nucleus. It may play a role in the postsynaptic responses of hippocampal neurons. This cytoplasmic protein is expressed in brain and testis. Defects in BRAF are involved in a wide range of cancers including lung cancer and non-Hodgkin lymphoma (NHL). This protein contains 1 zinc-dependent phorbol-ester and DAG binding domain.

**B-RAF Antibody (S445) Blocking peptide -  
References**

Hingorani, S.R., et al., Cancer Res. 63(17):5198-5202 (2003). Lee, J.W., et al., Br. J. Cancer 89(10):1958-1960 (2003). Davies, H., et al., Nature 417(6892):949-954 (2002). Naoki, K., et al., Cancer Res. 62(23):7001-7003 (2002). Stephens, R.M., et al., Mol. Cell. Biol. 12(9):3733-3742 (1992).

**Synonyms** BRAF1, RAFB1**Function**

Protein kinase involved in the transduction of mitogenic signals from the cell membrane to the nucleus (Probable). Phosphorylates MAP2K1, and thereby activates the MAP kinase signal transduction pathway (PubMed:<a href="http://www.uniprot.org/citations/21441910" target="\_blank">21441910</a>, PubMed:<a href="http://www.uniprot.org/citations/29433126" target="\_blank">29433126</a>). May play a role in the postsynaptic responses of hippocampal neurons (PubMed:<a href="http://www.uniprot.org/citations/1508179" target="\_blank">1508179</a>).

**Cellular Location**

Nucleus. Cytoplasm. Cell membrane. Note=Colocalizes with RGS14 and RAF1 in both the cytoplasm and membranes.

**Tissue Location**

Brain and testis.

**B-RAF Antibody (S445) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)