

TBB5 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP7858b**Specification****TBB5 Antibody (C-term) Blocking Peptide -
Product Information**Primary Accession [P07437](#)**TBB5 Antibody (C-term) Blocking Peptide -
Additional Information**

Gene ID 203068

Other NamesTubulin beta chain, Tubulin beta-5 chain,
TUBB, TUBB5**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP7858b](/products/AP7858b) was selected from the C-term region of human TBB5. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**TBB5 Antibody (C-term) Blocking Peptide -
Protein Information**

Name TUBB

Synonyms TUBB5

**TBB5 Antibody (C-term) Blocking Peptide -
Background**

Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain.

**TBB5 Antibody (C-term) Blocking Peptide -
References**

Wiesen,K.M., Cancer Lett. 257 (2), 227-235 (2007)Navarro-Nunez,L., Haematologica 92 (4), 513-518 (2007)Volz,A., Hum. Genet. 93 (1), 42-46 (1994)

Function

Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain.

Cellular Location

Cytoplasm, cytoskeleton

Tissue Location

Ubiquitously expressed with highest levels in spleen, thymus and immature brain.

TBB5 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)