

MEK2 (MAP2K2) Antibody (Center) Blocking peptide
Synthetic peptide
Catalog # BP7961c**Specification****MEK2 (MAP2K2) Antibody (Center) Blocking peptide - Product Information**Primary Accession [P36507](#)**MEK2 (MAP2K2) Antibody (Center) Blocking peptide - Additional Information****Gene ID** 5605**Other Names**

Dual specificity mitogen-activated protein kinase kinase 2, MAP kinase kinase 2, MAPKK 2, ERK activator kinase 2, MAPK/ERK kinase 2, MEK 2, MAP2K2, MEK2, MKK2, PRKMK2

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP7961c](/product/products/AP7961c) was selected from the Center region of human MAP2K2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

MEK2 (MAP2K2) Antibody (Center) Blocking peptide - Protein Information**MEK2 (MAP2K2) Antibody (Center) Blocking peptide - Background**

MAP2K2 is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. The inhibition or degradation of this kinase is found to be involved in the pathogenesis of Yersinia and anthrax.

MEK2 (MAP2K2) Antibody (Center) Blocking peptide - References

Burroughs, K.D., et al., Mol. Cancer Res. 1(4):312-322 (2003). Tran, H., et al., Mol. Cell. Biol. 23(20):7177-7188 (2003). Li, S.P., et al., Cancer Res. 63(13):3473-3477 (2003). Li, Y., et al., J. Biol. Chem. 278(16):13663-13671 (2003). Liu, X., et al., J. Biol. Chem. 277(42):39312-39319 (2002).

Name MAP2K2

Synonyms MEK2, MKK2, PRKMK2

Function

Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates the ERK1 and ERK2 MAP kinases (By similarity). Activates BRAF in a KSR1 or KSR2-dependent manner; by binding to KSR1 or KSR2 releases the inhibitory intramolecular interaction between KSR1 or KSR2 protein kinase and N-terminal domains which promotes KSR1 or KSR2-BRAF dimerization and BRAF activation (PubMed:29433126).

Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein. Note=Membrane localization is probably regulated by its interaction with KSR1.

**MEK2 (MAP2K2) Antibody (Center)
Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)