

MAP2K2 Antibody (T394) Blocking Peptide
Synthetic peptide
Catalog # BP7961d**Specification****MAP2K2 Antibody (T394) Blocking Peptide -
Product Information**Primary Accession [P36507](#)**MAP2K2 Antibody (T394) Blocking Peptide -
Additional Information****Gene ID** 5605**Other Names**

Dual specificity mitogen-activated protein kinase kinase 2, MAP kinase kinase 2, MAPKK 2, ERK activator kinase 2, MAPK/ERK kinase 2, MEK 2, MAP2K2, MEK2, MKK2, PRKMK2

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP7961d](/product/products/AP7961d) was selected from the T394 region of human MAP2K2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**MAP2K2 Antibody (T394) Blocking Peptide -
Protein Information****MAP2K2 Antibody (T394) Blocking Peptide
- Background**

MAP2K2 is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. The inhibition or degradation of this kinase is found to be involved in the pathogenesis of Yersinia and anthrax.

**MAP2K2 Antibody (T394) Blocking Peptide
- References**

Burroughs, K.D., et al., Mol. Cancer Res. 1(4):312-322 (2003). Tran, H., et al., Mol. Cell. Biol. 23(20):7177-7188 (2003). Li, S.P., et al., Cancer Res. 63(13):3473-3477 (2003). Li, Y., et al., J. Biol. Chem. 278(16):13663-13671 (2003). Liu, X., et al., J. Biol. Chem. 277(42):39312-39319 (2002).

Name MAP2K2

Synonyms MEK2, MKK2, PRKMK2

Function

Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates the ERK1 and ERK2 MAP kinases (By similarity). Activates BRAF in a KSR1 or KSR2-dependent manner; by binding to KSR1 or KSR2 releases the inhibitory intramolecular interaction between KSR1 or KSR2 protein kinase and N-terminal domains which promotes KSR1 or KSR2-BRAF dimerization and BRAF activation (PubMed:29433126).

Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein. Note=Membrane localization is probably regulated by its interaction with KSR1.

**MAP2K2 Antibody (T394) Blocking Peptide
- Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)