

STC1 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP7977c**Specification****STC1 Antibody (Center) Blocking Peptide -
Product Information**Primary Accession [P52823](#)**STC1 Antibody (Center) Blocking Peptide -
Additional Information****Gene ID** 6781**Other Names**

Stanniocalcin-1, STC-1, STC1, STC

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP7977c](/products/AP7977c) was selected from the Center region of human STC1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**STC1 Antibody (Center) Blocking Peptide -
Protein Information****Name** STC1**Synonyms** STC**STC1 Antibody (Center) Blocking Peptide -
Background**

STC1 is a secreted, homodimeric glycoprotein that is expressed in a wide variety of tissues and may have autocrine or paracrine functions. It contains 11 conserved cysteine residues and is phosphorylated by protein kinase C exclusively on its serine residues. This protein may play a role in the regulation of renal and intestinal calcium and phosphate transport, cell metabolism, or cellular calcium/phosphate homeostasis. Overexpression of human stanniocalcin 1 in mice produces high serum phosphate levels, dwarfism, and increased metabolic rate.

**STC1 Antibody (Center) Blocking Peptide -
References**

Law,A.Y., Exp. Cell Res. 314 (16), 2975-2984 (2008)
Holmes,D.I., Cell. Signal. 20 (3), 569-579 (2008)
Chang,A.C., Genomics 47 (3), 393-398 (1998)

Function

Stimulates renal phosphate reabsorption, and could therefore prevent hypercalcemia.

Cellular Location

Secreted.

Tissue Location

Expressed in most tissues, with the highest levels in ovary, prostate, heart, kidney and thyroid. In the kidney, expression is confined to the nephron, specifically in the distal convoluted tubule and in the collecting tubule. Not detected in the brain, liver, spleen, peripheral blood leukocytes and adrenal medulla

STC1 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)