

**PACSIN2 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP8088b****Specification****PACSIN2 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [Q9UNF0](#)**PACSIN2 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 11252**Other Names**

Protein kinase C and casein kinase substrate in neurons protein 2, Syndapin-2, Syndapin-II, PACSIN2

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP8088b](/product/products/AP8088b) was selected from the C-term region of human PACSIN2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**PACSIN2 Antibody (C-term) Blocking Peptide - Protein Information****Name** PACSIN2**PACSIN2 Antibody (C-term) Blocking Peptide - Background**

PACSIN may play a role in vesicle formation and transport. This protein homo- and hetero-aggregates with other PACSINs. It also binds dynamin 1, synaptojanin, synapsin 1 and the neural Wiskott-Aldrich syndrome protein (N-WASP). The protein exhibits a cytosolic-like cytoplasmic distribution and is ubiquitously expressed. PACSIN is phosphorylated by casein kinase 2 (CK2) and protein kinase C (PKC). The protein contains 1 FCH domain and 1 SH3 domain.

**PACSIN2 Antibody (C-term) Blocking Peptide - References**

Strausberg, R.L., et al., Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903 (2002). Wiemann, S., et al., Genome Res. 11(3):422-435 (2001). Ritter, B., et al., FEBS Lett. 454(3):356-362 (1999). Dunham, I., et al., Nature 402(6761):489-495 (1999).

**Function**

Regulates the morphogenesis and endocytosis of caveolae (By similarity). Lipid-binding protein that is able to promote the tubulation of the phosphatidic acid-containing membranes it preferentially binds. Plays a role in intracellular vesicle-mediated transport. Involved in the endocytosis of cell-surface receptors like the EGF receptor, contributing to its internalization in the absence of EGF stimulus.

**Cellular Location**

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasmic vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome. Recycling endosome membrane. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection. Membrane, caveola. Note=Detected at the neck of flask-shaped caveolae. Localization to tubular recycling endosomes probably requires interaction with MICALL1 and EHD1

**Tissue Location**

Widely expressed.

**PACSIN2 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)