

SKP2 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP8503c**Specification****SKP2 Antibody (Center) Blocking Peptide -
Product Information**Primary Accession [Q13309](#)**SKP2 Antibody (Center) Blocking Peptide -
Additional Information****Gene ID** 6502**Other Names**S-phase kinase-associated protein 2,
Cyclin-A/CDK2-associated protein p45,
F-box protein Skp2, F-box/LRR-repeat
protein 1, p45skp2, SKP2, FBXL1**Target/Specificity**

The synthetic peptide sequence used to generate the antibody AP8503c was selected from the Center region of human SKP2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**SKP2 Antibody (Center) Blocking Peptide -
Protein Information****Name** SKP2**SKP2 Antibody (Center) Blocking Peptide -
Background**

SKP2 is a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbls class; in addition to an F-box, this protein contains 10 tandem leucine-rich repeats. This protein is an essential element of the cyclin A-CDK2 S-phase kinase. It specifically recognizes phosphorylated cyclin-dependent kinase inhibitor 1B (CDKN1B, also referred to as p27 or KIP1) predominantly in S phase and interacts with S-phase kinase-associated protein 1 (SKP1 or p19). In addition, this gene is established as a protooncogene causally involved in the pathogenesis of lymphomas.

**SKP2 Antibody (Center) Blocking Peptide -
References**

Hussain,A.R., et.al., Leuk. Lymphoma 50 (7), 1204-1213 (2009)Yam,C.H., et.al., Mol. Cell. Biol. 19 (1), 635-645 (1999)

Synonyms FBXL1

Function

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription (PubMed:11931757, PubMed:12435635, PubMed:12769844, PubMed:12840033, PubMed:15342634, PubMed:15668399, PubMed:15949444, PubMed:16103164, PubMed:16262255, PubMed:16581786, PubMed:16951159, PubMed:17908926, PubMed:17962192, PubMed:22770219, PubMed:32267835). Specifically recognizes phosphorylated CDKN1B/p27kip and is involved in regulation of G1/S transition (By similarity).

Degradation of CDKN1B/p27kip also requires CKS1. Recognizes target proteins ORC1, CDT1, RBL2, KMT2A/MLL1, CDK9, RAG2, FOXO1, UBP43, YTHDF2, and probably MYC, TOB1 and TAL1 (PubMed:11931757, PubMed:12435635, PubMed:12769844, PubMed:12840033, PubMed:15342634, PubMed:15668399, PubMed:15949444, PubMed:16103164, PubMed:17962192, PubMed:16581786, PubMed:16951159, PubMed:17908926, PubMed:32267835).

Degradation of TAL1 also requires STUB1 (PubMed:17962192).

Recognizes CDKN1A in association with CCNE1 or CCNE2 and CDK2 (PubMed:16262255).

Promotes ubiquitination and destruction of CDH1 in a CK1-dependent manner, thereby regulating cell migration (PubMed:22770219).

Cellular Location

Cytoplasm. Nucleus

**SKP2 Antibody (Center) Blocking Peptide -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)