

**ADRA2B Antibody (Center) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP8566c****Specification****ADRA2B Antibody (Center) Blocking Peptide -  
Product Information**Primary Accession [P18089](#)**ADRA2B Antibody (Center) Blocking Peptide -  
Additional Information****Gene ID 151****Other Names**

Alpha-2B adrenergic receptor, Alpha-2  
adrenergic receptor subtype C2, Alpha-2B  
adrenoreceptor, Alpha-2B adrenoceptor,  
Alpha-2BAR, ADRA2B, ADRA2L1, ADRA2RL1

**Target/Specificity**

The synthetic peptide sequence used to  
generate the antibody <a  
href=/products/AP8566c>AP8566c</a>  
was selected from the Center region of  
human ADRA2B. A 10 to 100 fold molar  
excess to antibody is recommended.  
Precise conditions should be optimized for a  
particular assay.

**Format**

Peptides are lyophilized in a solid powder  
format. Peptides can be reconstituted in  
solution using the appropriate buffer as  
needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6  
months. For long term storage store at  
-20°C.

**Precautions**

This product is for research use only. Not  
for use in diagnostic or therapeutic  
procedures.

**ADRA2B Antibody (Center) Blocking Peptide -  
Protein Information****Name ADRA2B****ADRA2B Antibody (Center) Blocking  
Peptide - Background**

Alpha-2-adrenergic receptors are members of  
the G protein-coupled receptor superfamily.  
They include 3 highly homologous subtypes:  
alpha2A, alpha2B, and alpha2C. These  
receptors have a critical role in regulating  
neurotransmitter release from sympathetic  
nerves and from adrenergic neurons in the  
central nervous system. Alpha 2B adrenergic  
receptor subtype was observed to associate  
with eIF-2B, a guanine nucleotide exchange  
protein that functions in regulation of  
translation. A polymorphic variant of the  
alpha2B subtype, which lacks 3 glutamic acids  
from a glutamic acid repeat element, was  
identified to have decreased G protein-coupled  
receptor kinase-mediated phosphorylation and  
desensitization; this polymorphic form is also  
associated with reduced basal metabolic rate  
in obese subjects and may therefore contribute  
to the pathogenesis of obesity. Alpha 2B  
adrenergic receptor gene contains no introns  
in either its coding or untranslated sequences.

**ADRA2B Antibody (Center) Blocking  
Peptide - References**

Tabakoff,B., et.al., BMC Biol. 7, 70  
(2009) Weinshank,R.L., et.al. Mol. Pharmacol.  
38 (5), 681-688 (1990)

**Synonyms** ADRA2L1, ADRA2RL1**Function**

Alpha-2 adrenergic receptors mediate the catecholamine- induced inhibition of adenylate cyclase through the action of G proteins. The rank order of potency for agonists of this receptor is clonidine > norepinephrine > epinephrine = oxymetazoline > dopamine > p-tyramine = phenylephrine > serotonin > p-synephrine / p-octopamine. For antagonists, the rank order is yohimbine > chlorpromazine > phentolamine > mianserine > spiperone > prazosin > alprenolol > propanolol > pindolol.

**Cellular Location**

Cell membrane; Multi-pass membrane protein. Note=Interaction with RAB26, GGA1, GGA2 and GGA3 mediates transport from the Golgi to the cell membrane.

**ADRA2B Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)