

CBLB Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP8668c**Specification****CBLB Antibody (Center) Blocking Peptide -
Product Information**Primary Accession [Q13191](#)**CBLB Antibody (Center) Blocking Peptide -
Additional Information****Gene ID 868****Other Names**

E3 ubiquitin-protein ligase CBL-B, 632-,
Casitas B-lineage lymphoma
proto-oncogene b, RING finger protein 56,
SH3-binding protein CBL-B, Signal
transduction protein CBL-B, CBLB, RNF56

Target/Specificity

The synthetic peptide sequence used to
generate the antibody AP8668c
was selected from the Center region of
human CBLB. A 10 to 100 fold molar excess
to antibody is recommended. Precise
conditions should be optimized for a
particular assay.

Format

Peptides are lyophilized in a solid powder
format. Peptides can be reconstituted in
solution using the appropriate buffer as
needed.

Storage

Maintain refrigerated at 2-8°C for up to 6
months. For long term storage store at
-20°C.

Precautions

This product is for research use only. Not
for use in diagnostic or therapeutic
procedures.

**CBLB Antibody (Center) Blocking Peptide -
Protein Information****CBLB Antibody (Center) Blocking Peptide -
Background**

E3 ubiquitin-protein ligase which accepts
ubiquitin from specific E2 ubiquitin-conjugating
enzymes, and transfers it to substrates,
generally promoting their degradation by the
proteasome. Negatively regulates TCR (T-cell
receptor), BCR (B-cell receptor) and FCER1
(high affinity immunoglobulin epsilon receptor)
signal transduction pathways. In naive T-cells,
inhibits VAV1 activation upon TCR engagement
and imposes a requirement for CD28
costimulation for proliferation and IL-2
production. It also acts by promoting
PIK3R1/p85 ubiquitination, which impairs its
recruitment to the TCR and subsequent
activation. In activated T-cells, inhibits PLCG1
activation and calcium mobilization upon
restimulation and promotes anergy. In B-cells,
acts by ubiquitinating SYK and promoting its
proteasomal degradation. It may also be
involved in EGFR ubiquitination and
internalization.

**CBLB Antibody (Center) Blocking Peptide -
References**

Lavagna-Sevenier,C., et.al., J. Biol. Chem. 273
(24), 14962-14967 (1998)Yokoi,N., et.al.,
Biochem. Biophys. Res. Commun. 368 (1),
37-42 (2008)

Name CBLB

Synonyms RNF56

Function

E3 ubiquitin-protein ligase which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and transfers it to substrates, generally promoting their degradation by the proteasome. Negatively regulates TCR (T-cell receptor), BCR (B-cell receptor) and FCER1 (high affinity immunoglobulin epsilon receptor) signal transduction pathways. In naive T-cells, inhibits VAV1 activation upon TCR engagement and imposes a requirement for CD28 costimulation for proliferation and IL-2 production. Also acts by promoting PIK3R1/p85 ubiquitination, which impairs its recruitment to the TCR and subsequent activation. In activated T-cells, inhibits PLCG1 activation and calcium mobilization upon restimulation and promotes anergy. In B-cells, acts by ubiquitinating SYK and promoting its proteasomal degradation. Slightly promotes SRC ubiquitination. May be involved in EGFR ubiquitination and internalization. May be functionally coupled with the E2 ubiquitin-protein ligase UB2D3. In association with CBL, required for proper feedback inhibition of ciliary platelet-derived growth factor receptor-alpha (PDGFRA) signaling pathway via ubiquitination and internalization of PDGFRA (By similarity).

Cellular Location

Cytoplasm. Note=Upon EGF stimulation, associates with endocytic vesicles

Tissue Location

Expressed in placenta, heart, lung, kidney, spleen, ovary and testis, as well as fetal brain and liver and hematopoietic cell lines, but not in adult brain, liver, pancreas, salivary gland, or skeletal muscle. Present in lymphocytes (at protein level)

CBLB Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)