

MAP6 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP8670c**Specification****MAP6 Antibody (Center) Blocking Peptide -
Product Information**Primary Accession [Q96JE9](#)**MAP6 Antibody (Center) Blocking Peptide -
Additional Information****Gene ID** 4135**Other Names**Microtubule-associated protein 6, MAP-6,
Stable tubule-only polypeptide, STOP,
MAP6, KIAA1878**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP8670c](/products/AP8670c) was selected from the Center region of human MAP6. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**MAP6 Antibody (Center) Blocking Peptide -
Protein Information****Name** MAP6**MAP6 Antibody (Center) Blocking Peptide
- Background**

MAP6 is a microtubule-associated protein. This protein is a calmodulin-binding and calmodulin-regulated protein that is involved in microtubule stabilization.

**MAP6 Antibody (Center) Blocking Peptide
- References**

Andrieux,A., et.al., Genes Dev. 16 (18), 2350-2364 (2002)Bosc,C., et.al., Biochemistry 42 (42), 12125-12132 (2003)

Synonyms KIAA1878**Function**

Involved in microtubule stabilization in many cell types, including neuronal cells (By similarity). Specifically has microtubule cold stabilizing activity (By similarity). Involved in dendrite morphogenesis and maintenance by regulating lysosomal trafficking via its interaction with TMEM106B (PubMed: <http://www.uniprot.org/citations/24357581> target="_blank">24357581). Regulates KIF5A- mediated axonal cargo transport (By similarity). Regulates axonal growth during neuron polarization (By similarity).

Cellular Location

Cytoplasm, cytoskeleton. Golgi apparatus {ECO:0000250|UniProtKB:Q63560}. Cell projection, axon {ECO:0000250|UniProtKB:Q63560}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q63560}. Cytoplasmic vesicle, secretory vesicle membrane {ECO:0000250|UniProtKB:Q63560}; Lipid-anchor {ECO:0000250|UniProtKB:Q63560}; Cytoplasmic side {ECO:0000250|UniProtKB:Q63560}. Note=Localizes predominantly in the proximal part of the axon (By similarity). Preferentially is concentrated on a portion of the microtubule polymer in which tubulin is modified by detyrosination and acetylation and is also resistant to depolymerization induced by both nocodazole and cold (By similarity) In unpolarized neurons, localizes to the Golgi and to secretory vesicles accumulating transiently at the tips of a subset of neurites (By similarity). Following neuronal polarization and during axon outgrowth, accumulates in the axonal growth cone and subsequently localizes throughout the axon (By similarity). Partially localizes to dendrites in mature neurons (By similarity). Colocalizes with neurofilament (NF)-rich inclusions in spinal chord and brain neurons of patients with amyotrophic lateral sclerosis (ALS) (PubMed:14692697) {ECO:0000250|UniProtKB:Q63560, ECO:0000269|PubMed:14692697}

Tissue Location

Expressed in brain (at protein level).
Expressed in spinal cord. Isoform 2
expression is up-regulated in the prefrontal
cortex (Brodmann's area 46) of patients
with schizophrenia (postmortem brain
study).

MAP6 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Blocking Peptides](#)