

### **ALDH3B1 Antibody (Center) Blocking Peptide**

Synthetic peptide Catalog # BP8706c

### **Specification**

ALDH3B1 Antibody (Center) Blocking Peptide - Product Information

Primary Accession P43353

ALDH3B1 Antibody (Center) Blocking Peptide - Additional Information

Gene ID 221

#### **Other Names**

Aldehyde dehydrogenase family 3 member B1, Aldehyde dehydrogenase 7, ALDH3B1, ALDH7

### Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP8706c>AP8706c</a> was selected from the Center region of human ALDH3B1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ALDH3B1 Antibody (Center) Blocking Peptide - Protein Information

Name ALDH3B1

## ALDH3B1 Antibody (Center) Blocking Peptide - Background

The aldehyde dehydrogenases are a family of isozymes that may play a major role in the detoxification of aldehydes generated by alcohol metabolism and lipid peroxidation.

### ALDH3B1 Antibody (Center) Blocking Peptide - References

Marchitti, S.A., et.al., Biochem. Biophys. Res. Commun. 356 (3), 792-798 (2007)



### Synonyms ALDH7

#### **Function**

Oxidizes medium and long chain saturated and unsaturated aldehydes (PubMed: <a hre f="http://www.uniprot.org/citations/173822 92" target=" blank">17382292</a>, PubMed:<a href="http://www.uniprot.org/ci tations/23721920" target=" blank">23721920</a>). Metabolizes also benzaldehyde (PubMed:<a href="http://www.uniprot.org/citations/1738 2292" target=" blank">17382292</a>). Low activity towards acetaldehyde and 3,4-dihydroxyphenylacetaldehyde (PubMed:<a href="http://www.uniprot.org/c itations/17382292" target="\_blank">17382292</a>, PubMed:<a href="http://www.uniprot.org/ci tations/23721920" target=" blank">23721920</a>). May not metabolize short chain aldehydes. Can use both NADP(+) and NAD(+) as electron acceptor (PubMed:<a href="http://www.uni prot.org/citations/17382292" target=" blank">17382292</a>). May have a protective role against the cytotoxicity induced by lipid peroxidation (PubMed:<a href="http://www.uniprot.org/c itations/17382292" target=" blank">17382292</a>).

### **Cellular Location**

Cell membrane; Lipid-anchor. Note=Primarily in the plasma membrane as well as in some punctate structures in the cytoplasm

### **Tissue Location**

Highest expression in kidney and lung.

# ALDH3B1 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides