

RAC2 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP8839b**Specification****RAC2 Antibody (C-term) Blocking Peptide -
Product Information**Primary Accession [P15153](#)**RAC2 Antibody (C-term) Blocking Peptide -
Additional Information****Gene ID 5880****Other Names**Ras-related C3 botulinum toxin substrate 2,
GX, Small G protein, p21-Rac2, RAC2**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP8839b](/products/AP8839b) was selected from the C-term region of human RAC2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**RAC2 Antibody (C-term) Blocking Peptide -
Protein Information****Name RAC2****Function****RAC2 Antibody (C-term) Blocking Peptide -
Background**

RAC2 is a GTPase which belongs to the RAS superfamily of small GTP-binding proteins. Members of this superfamily appear to regulate a diverse array of cellular events, including the control of cell growth, cytoskeletal reorganization, and the activation of protein kinases.

**RAC2 Antibody (C-term) Blocking Peptide -
References**

Reibel,L., et.al., Biochem. Biophys. Res. Commun. 175 (2), 451-458 (1991) Brechard,S., et.al., J. Leukoc. Biol. 85 (4), 638-647 (2009)

Plasma membrane-associated small GTPase which cycles between an active GTP-bound and inactive GDP-bound state. In active state binds to a variety of effector proteins to regulate cellular responses, such as secretory processes, phagocytosis of apoptotic cells and epithelial cell polarization. Augments the production of reactive oxygen species (ROS) by NADPH oxidase.

Cellular Location

Cytoplasm. Note=Membrane-associated when activated

Tissue Location

Hematopoietic specific.

RAC2 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)