

**EED Antibody (Center) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP8896c****Specification****EED Antibody (Center) Blocking Peptide - Product Information**Primary Accession [075530](#)**EED Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 8726**Other Names**

Polycomb protein EED, hEED, WD protein associating with integrin cytoplasmic tails 1, WAIT-1, EED

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP8896c](/products/AP8896c) was selected from the Center region of human EED. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**EED Antibody (Center) Blocking Peptide - Protein Information****Name** EED ([HGNC:3188](#))**EED Antibody (Center) Blocking Peptide - Background**

EED is a polycomb group (PcG) protein. Component of the PRC2/EED-EZH2 complex, which methylates 'Lys-9' and 'Lys-27' of histone H3, leading to transcriptional repression of the affected target gene. The PRC2/EED-EZH2 complex may also serve as a recruiting platform for DNA methyltransferases, thereby linking two epigenetic repression systems.

**EED Antibody (Center) Blocking Peptide - References**

Dephoure N., et.al., Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).

**Function**

Polycomb group (PcG) protein. Component of the PRC2/EED-EZH2 complex, which methylates 'Lys-9' and 'Lys-27' of histone H3, leading to transcriptional repression of the affected target gene. Also recognizes 'Lys-26' trimethylated histone H1 with the effect of inhibiting PRC2 complex methyltransferase activity on nucleosomal histone H3 'Lys-27', whereas H3 'Lys-27' recognition has the opposite effect, enabling the propagation of this repressive mark. The PRC2/EED- EZH2 complex may also serve as a recruiting platform for DNA methyltransferases, thereby linking two epigenetic repression systems. Genes repressed by the PRC2/EED-EZH2 complex include HOXC8, HOXA9, MYT1 and CDKN2A.

**Cellular Location**

Nucleus. Chromosome. Note=Transiently colocalizes with XIST at inactive X chromosomes

**Tissue Location**

Expressed in brain, colon, heart, kidney, liver, lung, muscle, ovary, peripheral blood leukocytes, pancreas, placenta, prostate, spleen, small intestine, testis, thymus and uterus. Appears to be overexpressed in breast and colon cancer

**EED Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)