

CIRBP Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP9139b**Specification****CIRBP Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [Q14011](#)**CIRBP Antibody (C-term) Blocking Peptide - Additional Information****Gene ID 1153****Other Names**

Cold-inducible RNA-binding protein, A18 hnRNP, Glycine-rich RNA-binding protein CIRP, CIRBP, A18HNRNP, CIRP

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP9139b](#) was selected from the C-term region of human CIRBP. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CIRBP Antibody (C-term) Blocking Peptide - Protein Information**Name CIRBP****CIRBP Antibody (C-term) Blocking Peptide - Background**

CIRBP is cold-inducible mRNA binding protein that plays a protective role in the genotoxic stress response by stabilizing transcripts of genes involved in cell survival. It acts as a translational activator and seems to play an essential role in cold-induced suppression of cell proliferation. It binds specifically to the 3'-untranslated regions (3'-UTRs) of stress-responsive transcripts RPA2 and TXN. This protein acts as a translational repressor (By similarity) and promotes assembly of stress granules (SGs), when overexpressed.

CIRBP Antibody (C-term) Blocking Peptide - References

Colinge J., et.al., Submitted (OCT-2008) to UniProtKB. Yang R., et.al., Nucleic Acids Res. 34:1224-1236(2006).

Synonyms A18HNRNP, CIRP**Function**

Cold-inducible mRNA binding protein that plays a protective role in the genotoxic stress response by stabilizing transcripts of genes involved in cell survival. Acts as a translational activator. Seems to play an essential role in cold-induced suppression of cell proliferation. Binds specifically to the 3'-untranslated regions (3'-UTRs) of stress-responsive transcripts RPA2 and TXN. Acts as a translational repressor (By similarity). Promotes assembly of stress granules (SGs), when overexpressed.

Cellular Location

Nucleus, nucleoplasm. Cytoplasm
Note=Translocates from the nucleus to the cytoplasm after exposure to UV radiation. Translocates from the nucleus to the cytoplasm into stress granules upon various cytoplasmic stresses, such as osmotic and heat shocks. Its recruitment into stress granules occurs in the absence of TIAR proteins (By similarity).

Tissue Location

Ubiquitous.

**CIRBP Antibody (C-term) Blocking Peptide
- Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

CIRBP Antibody (C-term) Blocking Peptide - Citations

- [Cold-inducible RNA-binding protein mediates cold air inducible airway mucin production through TLR4/NF-κB signaling pathway.](#)