

Anti-ABCA4 (Rim Protein) Antibody

Our Anti-ABCA4 (Rim Protein) primary antibody from PhosphoSolutions is mouse monoclonal. It detects Catalog # AN1305

Specification

Anti-ABCA4 (Rim Protein) Antibody - Product Information

Application Primary Accession Reactivity Host Clonality

Isotype Calculated MW WB F1MWM0

Bovine, Chicken

Mouse Monoclonal IgG1 257376

Anti-ABCA4 (Rim Protein) Antibody - Additional Information

Gene ID **281584**

Other Names


ABC 10 antibody, ABC A4 antibody, ABC transporter retinal-specific antibody, ABC10 antibody, ABCA 4 antibody, abcA4 antibody, ABCA4_HUMAN antibody, ABCR antibody, ARMD 2 antibody, ARMD2 antibody, ATP binding cassette 10 antibody, ATP binding cassette sub family A member 4 antibody, ATP binding cassette transporter antibody, ATP binding cassette transporter retinal specific antibody, ATP binding cassette, sub family A (ABC1) member 4 antibody, ATP binding cassette sub family A (ABC1) member 4 antibody, ATP binding transporter retinal specific antibody, ATP-binding cassette sub-family A member 4 antibody, CORD 3 antibody, CORD3 antibody, DKFZp781N1972 antibody, FFM antibody, Retinal specific ATP binding cassette transporter antibody, Retinal specific ATP binding cassette transporter antibody, Retinal-specific ATP-binding cassette transporter antibody, RIM protein antibody, RmP antibody, RP 19 antibody, RP19 antibody, Stargardt disease protein antibody, STGD antibody, STGD1 antibody

Target/Specificity

ABCA4 (ATP-binding cassette, sub-family A (ABC1), member 4, Rim Protein) is a member of the superfamily of ATP-binding cassette (ABC) transporters (Illing et al., 1997). ABC proteins transport various molecules across extra- and intracellular membranes. This protein is a retina-specific ABC transporter with N-retinylidene-PE as a substrate. It is expressed exclusively in retina photoreceptor cells, indicating the gene product mediates transport of an essential molecule across the photoreceptor cell membrane. Mutations in this gene are found in patients diagnosed with Stargardt disease and are associated with retinitis pigmentosa-19 and age-related macular degeneration (Wiszniewski et al., 2003). Defects in ABCA4 are the cause of Stargardt disease type 1 (STGD1) (Molday et al., 2000). STGD is one of the most frequent causes of macular degeneration in childhood. Defects in ABCA4 are also known to cause fundus flavimaculatus (FFM), age-related macular degeneration type 2 (ARMD2) and cone-rod dystrophy type 3 (CORD3) (Klevering et al., 2005).

Dilution WB~~1:1000







Format

Protein G Purified

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-ABCA4 (Rim Protein) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Anti-ABCA4 (Rim Protein) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-ABCA4 (Rim Protein) Antibody - Images



Immunohistochemical staining of adult mouse retina showing specific immunolabeling of the ABCA4 protein (red, 1;100). Photo courtesy of Mary Raven, University of California, Santa Barbara, CA.

Anti-ABCA4 (Rim Protein) Antibody - Background

ABCA4 (ATP-binding cassette, sub-family A (ABC1), member 4, Rim Protein) is a member of the superfamily of ATP-binding cassette (ABC) transporters (Illing et al., 1997). ABC proteins transport various molecules across extra- and intracellular membranes. This protein is a retina-specific ABC transporter with N-retinylidene-PE as a substrate. It is expressed exclusively in retina photoreceptor cells, indicating the gene product mediates transport of an essential molecule across the photoreceptor cell membrane. Mutations in this gene are found in patients diagnosed with Stargardt disease and are associated with retinitis pigmentosa-19 and age-related macular





Tel: 858.875.1900 Fax: 858.875.1999

degeneration (Wiszniewski et al., 2003). Defects in ABCA4 are the cause of Stargardt disease type 1 (STGD1) (Molday et al., 2000). STGD is one of the most frequent causes of macular degeneration in childhood. Defects in ABCA4 are also known to cause fundus flavimaculatus (FFM), age-related macular degeneration type 2 (ARMD2) and cone-rod dystrophy type 3 (CORD3) (Klevering et al., 2005).