

Anti-MEF2C (Ser222) Antibody

Our Anti-MEF2C (Ser222) rabbit polyclonal phosphospecific primary antibody from PhosphoSolutions is
Catalog # AN1446

Specification

Anti-MEF2C (Ser222) Antibody - Product Information

Primary Accession	Q06413
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	51221

Anti-MEF2C (Ser222) Antibody - Additional Information

Gene ID **4208**

Other Names

C5DELq14.3 antibody, DEL5q14.3 antibody, MADS box transcription enhancer factor 2 polypeptide C (myocyte enhancer factor 2C) antibody, MADS box transcription enhancer factor 2 polypeptide C antibody, MEF2C antibody, MEF2C_HUMAN antibody, Myocyte enhancer factor 2C antibody, Myocyte specific enhancer factor 2C antibody, Myocyte-specific enhancer factor 2C antibody, OTTHUMP00000222409 antibody, Similar to MADS box transcription enhancer factor 2 polypeptide C antibody

Target/Specificity

MEF2C, also known as MADS box transcription enhancer factor 2, polypeptide C, is one of 4 MEF2 (myocyte enhancer factor 2) transcription factors that encode proteins for development of skeletal muscle and brain proliferation and differentiation (McDermott et al, 1993), along with regulating stress-response during cardiac hypertrophy in mammals (Wu et al, 2015). Phosphorylation of MEF2C at serine 59 has been shown to be negatively regulated by integrin-linked kinase (ILK) (Dong et al, 2015). Recently, high MEF2C expression has been associated with a subset of acute myeloid leukemia (AML) patients with adverse-risk disease features and poor outcomes (Laszlo et al, 2015). Phosphorylation of MEF2C at serine 222 may play a key role in MEF2C signaling.

Format

Antigen Affinity Purified from Pooled Serum

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-MEF2C (Ser222) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Anti-MEF2C (Ser222) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-MEF2C (Ser222) Antibody - Images



Western blot of OCIAML2 lysate showing specific immunolabeling of the ~51 kDa MEF2C phosphorylated at Ser222 in the first lane (-). Phosphospecificity is shown in the second lane (+) where the immunolabeling is completely eliminated by blot treatment with lambda phosphatase ((λ-Ptase, 1200 units for 30 minutes).

Anti-MEF2C (Ser222) Antibody - Background

MEF2C, also known as MADS box transcription enhancer factor 2, polypeptide C, is one of 4 MEF2 (myocyte enhancer factor 2) transcription factors that encode proteins for development of skeletal muscle and brain proliferation and differentiation (McDermott et al, 1993), along with regulating stress-response during cardiac hypertrophy in mammals (Wu et al, 2015). Phosphorylation of MEF2C at serine 59 has been shown to be negatively regulated by integrin-linked kinase (ILK) (Dong et al, 2015). Recently, high MEF2C expression has been associated with a subset of acute myeloid leukemia (AML) patients with adverse-risk disease features and poor outcomes (Laszlo et al, 2015). Phosphorylation of MEF2C at serine 222 may play a key role in MEF2C signaling.