

Anti-Arpc1b/p41-Arc (C-terminal region) Antibody
Catalog # AN1642**Specification****Anti-Arpc1b/p41-Arc (C-terminal region) Antibody - Product Information**

Application	WB, IHC
Primary Accession	O15143
Reactivity	Bovine, Chicken
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	40950

Anti-Arpc1b/p41-Arc (C-terminal region) Antibody - Additional Information

Gene ID	10095
Other Names	
Arc41, p41, Arpc1b	

Target/Specificity

Cellular morphology, adhesion, and motility occur through dynamic reorganization of actin-based superstructures. Actin-binding proteins are critical for regulating actin polymerization and superstructure formation. The Arp2/3 complex is an actin polymerization-inducing complex that includes Arp2, Arp3, p41-Arc, p34-Arc, p21-Arc, p20-Arc, and p16-Arc. Several nucleation promoting factors, such as WASP and coronin, regulate the activity of the Arp2/3 complex. In addition, the Arp2/3 complex may be regulated by phosphorylation of specific subunits in the complex. p41-Arc (Arpc1b) subunit Arpc1 has two isoforms in humans, Arpc1a and Arpc1b. PAK1 can bind and phosphorylate Thr-21 in Arpc1b leading to growth factor-stimulated cell motility. In addition, Arpc1b colocalizes with γ -tubulin at centrosomes and stimulates Aurora A activity. Aurora A phosphorylates Arpc1b on Thr-21 and a nonphosphorylatable Arpc1b mutant cannot activate Aurora A kinase and centrosome amplification. Thus, Arpc1b has roles in cytoskeletal dynamics during cell motility and mitosis, and these activities are regulated by phosphorylation at Thr-21

Dilution

WB~~1:1000
IHC~~1:100~500

Format

Antigen Affinity Purified

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Arpc1b/p41-Arc (C-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Anti-Arpc1b/p41-Arc (C-terminal region) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Arpc1b/p41-Arc (C-terminal region) Antibody - Images



Western blot of rat A7r5 (lane 1), human Jurkat (lane 2), rat PC12 (lane 3), human PC3 (lane 4), mouse C2C12 (lane 5), and human A431 cells (lane 6). The blot was probed with rabbit polyclonal anti-Arpc1b (C-terminal region).

Anti-Arpc1b/p41-Arc (C-terminal region) Antibody - Background

Cellular morphology, adhesion, and motility occur through dynamic reorganization of actin-based superstructures. Actin-binding proteins are critical for regulating actin polymerization and superstructure formation. The Arp2/3 complex is an actin polymerization-inducing complex that includes Arp2, Arp3, p41-Arc, p34-Arc, p21-Arc, p20-Arc, and p16-Arc. Several nucleation promoting factors, such as WASP and coronin, regulate the activity of the Arp2/3 complex. In addition, the Arp2/3 complex may be regulated by phosphorylation of specific subunits in the complex. p41-Arc (Arpc1b) subunit Arpc1 has two isoforms in humans, Arpc1a and Arpc1b. PAK1 can bind and phosphorylate Thr-21 in Arpc1b leading to growth factor-stimulated cell motility. In addition, Arpc1b colocalizes with γ -tubulin at centrosomes and stimulates Aurora A activity. Aurora A phosphorylates Arpc1b on Thr-21 and a nonphosphorylatable Arpc1b mutant cannot activate Aurora A kinase and centrosome amplification. Thus, Arpc1b has roles in cytoskeletal dynamics during cell motility and mitosis, and these activities are regulated by phosphorylation at Thr-21