

Anti-DAAM1 (N-terminal region) Antibody
Catalog # AN1739**Specification**

Anti-DAAM1 (N-terminal region) Antibody - Product Information

Application	WB
Primary Accession	Q9Y4D1
Reactivity	Bovine
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG2a
Calculated MW	123473

Anti-DAAM1 (N-terminal region) Antibody - Additional Information

Gene ID 23002

Other Names

formin, Disheveled morphogenesis

Target/Specificity

Formins include several families of proteins that regulate actin cytoskeletal dynamics via two conserved formin homology domains, FH1 and FH2. The FH1 region contains poly-proline stretches that promote interactions with profilin. The FH2 domain, located C-terminally to the FH1 domain, is highly conserved in formin proteins and possesses actin nucleation and polymerization activities. Through cooperation of FH1 and FH2, formins construct actin-based structures comprising linear, unbranched filaments that are used in stress fibers, actin cables, microspikes, and contractile rings. Dishevelled associated activator of morphogenesis proteins (DAAM1 and DAAM2) are formin family members involved in WNT signaling. DAAM1 is ubiquitously expressed and may be important for regulating actin dynamics in several cell types. DAAM1 can bind RhoGTPase and dishevelled in WNT signaling pathways, and interacts with the SH3 domains of cell signaling mediators, such as c-Src.

Dilution

WB~~1:1000

Format

Antigen Affinity Purified

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-DAAM1 (N-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Anti-DAAM1 (N-terminal region) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-DAAM1 (N-terminal region) Antibody - Images



Immunocytochemical labeling of DAAM1 in aldehyde-fixed and NP-40-permeabilized A431 cells. The cells were labeled with mouse monoclonal DAAM1 (N-terminal region) antibody, then the antibody was detected using appropriate secondary antibody conjugated to DyLight® 594. The corresponding phase image is shown to the right.

Anti-DAAM1 (N-terminal region) Antibody - Background

Formins include several families of proteins that regulate actin cytoskeletal dynamics via two conserved formin homology domains, FH1 and FH2. The FH1 region contains poly-proline stretches that promote interactions with profilin. The FH2 domain, located C-terminally to the FH1 domain, is highly conserved in formin proteins and possesses actin nucleation and polymerization activities. Through cooperation of FH1 and FH2, formins construct actin-based structures comprising linear, unbranched filaments that are used in stress fibers, actin cables, microspikes, and contractile rings. Dishevelled associated activator of morphogenesis proteins (DAAM1 and DAAM2) are formin family members involved in WNT signaling. DAAM1 is ubiquitously expressed and may be important for regulating actin dynamics in several cell types. DAAM1 can bind RhoGTPase and dishevelled in WNT signaling pathways, and interacts with the SH3 domains of cell signaling mediators, such as c-Src.