

Anti-MuRF1 (C-terminal region) Antibody
Catalog # AN1842**Specification****Anti-MuRF1 (C-terminal region) Antibody - Product Information**

Application	WB
Primary Accession	Q969Q1
Reactivity	Bovine
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	40248

Anti-MuRF1 (C-terminal region) Antibody - Additional InformationGene ID **84676****Other Names**

Tripartite motif 63, TRIM63, SMRZ, IRF, RNF28, Muscle RING finger, MURF-1

Target/Specificity

Muscle proteolysis is regulated by the ATP-dependent ubiquitin-proteasome system. This system involves ubiquitination of specific proteins, leading to recognition and degradation by the 26S proteasome complex. Ubiquitination requires interactions with ubiquitin related proteins, ubiquitin-activating (E1), ubiquitin-conjugating (E2) and ubiquitin-ligating enzymes (E3) known as ligases. Two muscle specific ubiquitin ligases have been identified, muscle ring finger 1 (MuRF-1) and Atrogin 1. Both ligases are regulated by the Akt1/FOXO1 signaling pathway, and both proteins have been shown to be upregulated prior to the onset of atrophy in multiple models of muscle wasting, including disuse and cachexia. MuRF1 is also known as TRIM63, SMRZ, and RNF28, and its expression is upregulated after TNF α treatment in C2C12 cells and muscle tissue, while localization of MuRF1 protein has been observed in the cytoplasm and nucleus of cells.

Dilution

WB~~1:1000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-MuRF1 (C-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Anti-MuRF1 (C-terminal region) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-MuRF1 (C-terminal region) Antibody - Images



Western blot analysis of mouse heart tissue (lanes 1 & 3) or C2C12 cells (lanes 2 & 4). The blot was probed with anti-MuRF1 (C-terminal region) (lanes 1 & 2) or anti-Atrogin-1 (lanes 3 & 4).



Immunocytochemical labeling of MuRF1 in mouse C2C12 cells. The cells were labeled with rabbit polyclonal MuRF1 antibody, then detected using appropriate secondary antibody conjugated to Cy3. The antibody was used in the absence (left) or presence (right) of blocking peptide (MX3405).

Anti-MuRF1 (C-terminal region) Antibody - Background

Muscle proteolysis is regulated by the ATP-dependent ubiquitin-proteasome system. This system involves ubiquitination of specific proteins, leading to recognition and degradation by the 26S proteasome complex. Ubiquitination requires interactions with ubiquitin related proteins, ubiquitin-activating (E1), ubiquitin-conjugating (E2) and ubiquitin-ligating enzymes (E3) known as ligases. Two muscle specific ubiquitin ligases have been identified, muscle ring finger 1 (MuRF-1) and Atrogin 1. Both ligases are regulated by the Akt1/FOXO1 signaling pathway, and both proteins have been shown to be upregulated prior to the onset of atrophy in multiple models of muscle wasting, including disuse and cachexia. MuRF1 is also known as TRIM63, SMRZ, and RNF28, and its expression is upregulated after TNF α treatment in C2C12 cells and muscle tissue, while localization of MuRF1 protein has been observed in the cytoplasm and nucleus of cells.