

CDH12 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP1473a

Specification

CDH12 Antibody (N-term) - Product Information

Application WB, IHC-P,E **Primary Accession** P55289 Other Accession Q5RIH3 Reactivity Human Predicted Mouse Host Rabbit Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 88332 Antigen Region 62-90

CDH12 Antibody (N-term) - Additional Information

Gene ID 1010

Other Names

Cadherin-12, Brain cadherin, BR-cadherin, Neural type cadherin 2, N-cadherin 2, CDH12

Target/Specificity

This CDH12 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 62-90 amino acids from the N-terminal region of human CDH12.

Dilution

WB~~1:1000 IHC-P~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDH12 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CDH12 Antibody (N-term) - Protein Information

Name CDH12





Function Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types.

Cellular Location

Cell membrane; Single-pass type I membrane protein


Tissue Location Brain.

CDH12 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CDH12 Antibody (N-term) - Images



Western blot analysis of anti-CDH12 Antibody (N-term) (RB13742) in Jurkat cell line lysates (35ug/lane). CDH12(arrow) was detected using the purified Pab.





Formalin-fixed and paraffin-embedded human brain tissue reacted with CDH12 antibody (N-term) (Cat.#AP1473a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

CDH12 Antibody (N-term) - Background

CDH12 is a type II classical cadherin from the cadherin superfamily of integral membrane proteins that mediate calcium-dependent cell-cell adhesion. Mature cadherin proteins are composed of a large N-terminal extracellular domain, a single membrane-spanning domain, and a small, highly conserved C-terminal cytoplasmic domain. Type II (atypical) cadherins are defined based on their lack of a HAV cell adhesion recognition sequence specific to type I cadherins. This particular cadherin appears to be expressed specifically in the brain and its temporal pattern of expression would be consistent with a role during a critical period of neuronal development, perhaps specifically during synaptogenesis.

CDH12 Antibody (N-term) - References

Shimoyama, Y., Biochem. J. 349 (PT 1), 159-167 (2000) Chalmers, I.J., Genomics 57 (1), 160-163 (1999) Kremmidiotis, G., Genomics 49 (3), 467-471 (1998) Selig, S., Proc. Natl. Acad. Sci. U.S.A. 94 (6), 2398-2403 (1997)