

**WWTR1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP16419b****Specification**

---

**WWTR1 Antibody (C-term) - Product Information**

|                   |                                                                                       |
|-------------------|---------------------------------------------------------------------------------------|
| Application       | WB,E                                                                                  |
| Primary Accession | <a href="#">O9GZV5</a>                                                                |
| Other Accession   | <a href="#">O9EPK5</a> , <a href="#">NP_001161752.1</a> , <a href="#">NP_056287.1</a> |
| Reactivity        | Human                                                                                 |
| Predicted         | Mouse                                                                                 |
| Host              | Rabbit                                                                                |
| Clonality         | Polyclonal                                                                            |
| Isotype           | Rabbit IgG                                                                            |
| Calculated MW     | 44101                                                                                 |
| Antigen Region    | 370-398                                                                               |

**WWTR1 Antibody (C-term) - Additional Information****Gene ID** 25937**Other Names**

WW domain-containing transcription regulator protein 1, Transcriptional coactivator with PDZ-binding motif, WWTR1, TAZ

**Target/Specificity**

This WWTR1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 370-398 amino acids from the C-terminal region of human WWTR1.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

WWTR1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**WWTR1 Antibody (C-term) - Protein Information****Name** WWTR1 ([HGNC:24042](#))

**Function** Transcriptional coactivator which acts as a downstream regulatory target in the Hippo signaling pathway that plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis (PubMed:[11118213](#), PubMed:[18227151](#), PubMed:[23911299](#)). The core of this pathway is composed of a kinase cascade wherein STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ (PubMed:[18227151](#)). WWTR1 enhances PAX8 and NKX2-1/TTF1-dependent gene activation (PubMed:[19010321](#)). In conjunction with YAP1, involved in the regulation of TGFβ1-dependent SMAD2 and SMAD3 nuclear accumulation (PubMed:[18568018](#)). Plays a key role in coupling SMADs to the transcriptional machinery such as the mediator complex (PubMed:[18568018](#)). Regulates embryonic stem-cell self-renewal, promotes cell proliferation and epithelial-mesenchymal transition (PubMed:[18227151](#), PubMed:[18568018](#)).

#### Cellular Location

Nucleus. Cytoplasm. Cell membrane. Cell junction, tight junction  
{ECO:0000250|UniProtKB:A0A8I3PQN6}. Note=Concentrates along specific portions of the plasma membrane, and accumulates in punctate nuclear bodies (By similarity). When phosphorylated, is retained in the cytoplasm by YWHAZ (By similarity). Can be retained in the nucleus by MED15 (PubMed:[18568018](#)). Localized in the cytoplasm in areas of epithelial cell high density (PubMed:[21145499](#)). At blastocyst stage expressed in the nucleus in trophectodermal cells, however expressed in the cytoplasm in the inner cell mass (By similarity). In the nucleus, phosphorylation by PRP4K induces nuclear exclusion (PubMed:[29695716](#)) Interaction with AMOTL2 results in localization to the cytoplasm and tight junctions (PubMed:[23911299](#)).  
{ECO:0000250|UniProtKB:Q9EPK5, ECO:0000269|PubMed:[18568018](#), ECO:0000269|PubMed:[21145499](#), ECO:0000269|PubMed:[23911299](#), ECO:0000269|PubMed:[29695716](#)}

#### Tissue Location


Highly expressed in kidney, heart, placenta and lung. Expressed in the thyroid tissue.


#### WWTR1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### WWTR1 Antibody (C-term) - Images





WWTR1 Antibody (C-term) (Cat. #AP16419b) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the WWTR1 antibody detected the WWTR1 protein (arrow).

#### **WWTR1 Antibody (C-term) - Background**

Transcriptional coactivator which acts as a downstream regulatory target in the Hippo signaling pathway that plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein MST1/MST2, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. WWTR1 enhances PAX8 and NKX2-1/TTF1-dependent gene activation. Regulates the nuclear accumulation of SMADS and has a key role in coupling them to the transcriptional machinery such as the mediator complex. Regulates embryonic stem-cell self-renewal, promotes cell proliferation and epithelial-mesenchymal transition.

#### **WWTR1 Antibody (C-term) - References**

- Remue, E., et al. FEBS Lett. 584(19):4175-4180(2010)
- Strakova, Z., et al. Biol. Reprod. 82(6):1112-1118(2010)
- Cho, H.H., et al. J. Cell. Physiol. 223(1):168-177(2010)
- McGeachie, M., et al. Circulation 120(24):2448-2454(2009)
- Ferrara, A.M., et al. J. Endocrinol. Invest. 32(3):238-241(2009)