

NRIP2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17224C**Specification**

NRIP2 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O9BQI9
Other Accession	NP_113662.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	31331
Antigen Region	91-119

NRIP2 Antibody (Center) - Additional Information**Gene ID** 83714**Other Names**

Nuclear receptor-interacting protein 2, NRIP2

Target/Specificity

This NRIP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 91-119 amino acids from the Central region of human NRIP2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions


NRIP2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NRIP2 Antibody (Center) - Protein Information**Name** NRIP2**Function** Down-regulates transcriptional activation by nuclear receptors such as NR1F2.


Cellular Location
Nucleus.**NRIP2 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NRIP2 Antibody (Center) - Images

NRIP2 Antibody (Center) (Cat. #AP17224c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the NRIP2 antibody detected the NRIP2 protein (arrow).



Anti-NRIP2 Antibody (Center) at 1:1000 dilution + human kidney lysate Lysates/proteins at 20 µg

per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 31 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

NRIP2 Antibody (Center) - Background

Down-regulates transcriptional activation by nuclear receptors such as NR1F2 (By similarity).

NRIP2 Antibody (Center) - References

Nakashima, K., et al. Allergol Int 55(1):77-83(2006)
Simpson, J.C., et al. EMBO Rep. 1(3):287-292(2000)
Greiner, E.F., et al. Proc. Natl. Acad. Sci. U.S.A. 97(13):7160-7165(2000)