

**DNAJC5G Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP17525c****Specification**

---

**DNAJC5G Antibody (Center) - Product Information**

|                   |                             |
|-------------------|-----------------------------|
| Application       | WB,E                        |
| Primary Accession | <a href="#">Q8N7S2</a>      |
| Other Accession   | <a href="#">NP_775921.1</a> |
| Reactivity        | Human                       |
| Host              | Rabbit                      |
| Clonality         | Polyclonal                  |
| Isotype           | Rabbit IgG                  |
| Calculated MW     | 21433                       |
| Antigen Region    | 88-115                      |

**DNAJC5G Antibody (Center) - Additional Information****Gene ID** 285126**Other Names**

DnaJ homolog subfamily C member 5G, Cysteine string protein-gamma, CSP-gamma, DNAJC5G

**Target/Specificity**

This DNAJC5G antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 88-115 amino acids from the Central region of human DNAJC5G.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

DNAJC5G Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**DNAJC5G Antibody (Center) - Protein Information****Name** DNAJC5G**Cellular Location**

Membrane; Lipid-anchor

**Tissue Location**


Testis specific.

**DNAJC5G Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**DNAJC5G Antibody (Center) - Images**



Anti-DNAJC5G Antibody (Center) at 1:1000 dilution + NCI-H292 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 21 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

**DNAJC5G Antibody (Center) - Background**

DNAJC5G contains 1 J domain.

**DNAJC5G Antibody (Center) - References**

Gerhard, D.S., et al. Genome Res. 14 (10B), 2121-2127 (2004) :