

**NTAN1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP17739b****Specification**

---

**NTAN1 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O96AB6</a>
Other Accession	<a href="#">NP_775745.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	34677
Antigen Region	203-230

**NTAN1 Antibody (C-term) - Additional Information****Gene ID** 123803**Other Names**

Protein N-terminal asparagine amidohydrolase, 351-, Protein NH2-terminal asparagine amidohydrolase, PNAA, Protein NH2-terminal asparagine deamidase, PNAD, Protein N-terminal Asn amidase, Protein N-terminal asparagine amidase, Protein NTN-amidase, NTAN1

**Target/Specificity**

This NTAN1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 203-230 amino acids from the C-terminal region of human NTAN1.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NTAN1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**NTAN1 Antibody (C-term) - Protein Information****Name** NTAN1

**Function** N-terminal asparagine deamidase that mediates deamidation of N-terminal asparagine residues to aspartate. Required for the ubiquitin-dependent turnover of intracellular proteins that initiate with Met-Asn. These proteins are acetylated on the retained initiator methionine and can subsequently be modified by the removal of N-acetyl methionine by acylaminoacid hydrolase (AAH). Conversion of the resulting N-terminal asparagine to aspartate by NTAN1/PNAD renders the protein susceptible to arginylation, polyubiquitination and degradation as specified by the N-end rule. This enzyme does not act on substrates with internal or C-terminal asparagines and does not act on glutamine residues in any position, nor on acetylated N-terminal peptidyl Asn.

#### Cellular Location


Cytoplasm {ECO:0000250|UniProtKB:Q28955}.

### NTAN1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### NTAN1 Antibody (C-term) - Images



NTAN1 Antibody (C-term) (Cat. #AP17739b) western blot analysis in T47D cell line lysates (35ug/lane). This demonstrates the NTAN1 antibody detected the NTAN1 protein (arrow).

### NTAN1 Antibody (C-term) - Background

Side-chain deamidation of N-terminal asparagine residues to aspartate. Required for the ubiquitin-dependent turnover of intracellular proteins that initiate with Met-Asn. These proteins are acetylated on the retained initiator methionine and can subsequently be modified by the removal of N-acetyl methionine by acylaminoacid hydrolase (AAH). Conversion of the resulting N-terminal asparagine to aspartate by PNAD renders the protein susceptible to arginylation, polyubiquitination and degradation as specified by the N-end rule. This enzyme does not act on substrates with internal or C-terminal asparagines and does not act on glutamine residues in any position (By similarity).

**NTAN1 Antibody (C-term) - References**

Okada, Y., et al. Hum. Mol. Genet. 19(11):2303-2312(2010)  
Kamdem, L.K., et al. Pharmacogenet. Genomics 18(6):507-514(2008)  
Lamesch, P., et al. Genomics 89(3):307-315(2007)  
Grigoryev, S., et al. J. Biol. Chem. 271(45):28521-28532(1996)