

MME Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20696a**Specification**

MME Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P08473
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	85514

MME Antibody (N-term) - Additional Information**Gene ID** 4311**Other Names**

Neprilysin, Atriopeptidase, Common acute lymphocytic leukemia antigen, CALLA, Enkephalinase, Neutral endopeptidase 2411, NEP, Neutral endopeptidase, Skin fibroblast elastase, SFE, CD10, MME, EPN

Target/Specificity

This MME antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 99-132 amino acids from the N-terminal region of human MME.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MME Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MME Antibody (N-term) - Protein Information**Name** MME {ECO:0000303|PubMed:27588448, ECO:0000312|HGNC:HGNC:7154}**Function** Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30

amino acids (PubMed:[15283675](#), PubMed:[6208535](#), PubMed:[6349683](#), PubMed:[8168535](#)). Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by cleavage of a Gly-Phe bond (PubMed:[17101991](#), PubMed:[6349683](#)). Catalyzes cleavage of bradykinin, substance P and neurotensin peptides (PubMed:[6208535](#)). Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9 (PubMed:[15283675](#), PubMed:[6349683](#)). Involved in the degradation of atrial natriuretic factor (ANF) and brain natriuretic factor (BNP(1-32)) (PubMed:[16254193](#), PubMed:[2531377](#), PubMed:[2972276](#)). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers (PubMed:[20876573](#)).

Cellular Location


Cell membrane; Single-pass type II membrane protein

MME Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MME Antibody (N-term) - Images



Western blot analysis of lysates from LNCaP, M. kidney, R. kidney cell line (from left to right), using MME Antibody (N-term)(Cat. #AP20696a). AP20696a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

MME Antibody (N-term) - Background

Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino acids. Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by cleavage of a Gly-Phe bond. Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9.

Involved in the degradation of atrial natriuretic factor (ANF). Displays UV- inducible elastase activity toward skin preelastic and elastic fibers.

MME Antibody (N-term) - References

Letarte M.,et al.J. Exp. Med. 168:1247-1253(1988).
Shipp M.A.,et al.Proc. Natl. Acad. Sci. U.S.A. 85:4819-4823(1988).
D'Adamio L.,et al.Proc. Natl. Acad. Sci. U.S.A. 86:7103-7107(1989).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.