

## **DVL1L1 Antibody (N-Term)**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21836a

## **Specification**

# **DVL1L1 Antibody (N-Term) - Product Information**

Application WB.E **Primary Accession** P54792 Reactivity Human Host **Rabbit** Clonality polyclonal Isotype Rabbit IgG Calculated MW 73254 Antigen Region 39-71

## **DVL1L1 Antibody (N-Term) - Additional Information**

#### **Other Names**

Putative segment polarity protein dishevelled homolog DVL1P1, DSH homolog 1-like, Segment polarity protein dishevelled homolog DVL-1-like, Dishevelled-1-like, DVL1P1, DVL, DVL1, DVL1L1

### Target/Specificity

This DVL1L1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 39-71 amino acids from human DVL1L1.

### **Dilution**

WB~~1:2000

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

# **Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

DVL1L1 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## **DVL1L1 Antibody (N-Term) - Protein Information**

### Name DVL1P1

Synonyms DVL, DVL1, DVL1L1

Function May play a role in the signal transduction pathway mediated by multiple Wnt genes.



**Cellular Location** Cytoplasm.

#### **Tissue Location**


Expressed in thymus, heart, liver, kidney, brain, skeletal muscle, and pancreas.

# **DVL1L1 Antibody (N-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **DVL1L1 Antibody (N-Term) - Images**



Anti-DVL1L1 Antibody (N-Term) at 1:2000 dilution + human heart lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 73 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

# **DVL1L1 Antibody (N-Term) - Background**

May play a role in the signal transduction pathway mediated by multiple Wnt genes.

## **DVL1L1 Antibody (N-Term) - References**

Pizzuti A., et al. Am. J. Hum. Genet. 58:722-729(1996).