

VLDLR Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21837b**Specification**

VLDLR Antibody (C-Term) - Product Information

Application	WB,E
Primary Accession	P98155
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	96098
Antigen Region	608-642

VLDLR Antibody (C-Term) - Additional Information**Gene ID** 7436**Other Names**

Very low-density lipoprotein receptor, VLDL receptor, VLDL-R, VLDLR

Target/Specificity

This VLDLR antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 608-642 amino acids from human VLDLR.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

VLDLR Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

VLDLR Antibody (C-Term) - Protein Information**Name** VLDLR

Function Multifunctional cell surface receptor that binds VLDL and transports it into cells by endocytosis and therefore plays an important role in energy metabolism. Also binds to a wide

range of other molecules including Reelin/RELN or apolipoprotein E/APOE- containing ligands as well as clusterin/CLU (PubMed:[24381170](#), PubMed:[30873003](#)). In the off-state of the pathway, forms homooligomers or heterooligomers with LRP8 (PubMed:[30873003](#)). Upon binding to ligands, homooligomers are rearranged to higher order receptor clusters that transmit the extracellular RELN signal to intracellular signaling processes by binding to DAB1 (PubMed:[30873003](#)). This interaction results in phosphorylation of DAB1 leading to the ultimate cell responses required for the correct positioning of newly generated neurons. Later, mediates a stop signal for migrating neurons, preventing them from entering the marginal zone (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein Membrane, clathrin-coated pit; Single-pass type I membrane protein

Tissue Location


Abundant in heart and skeletal muscle; also ovary and kidney; not in liver

VLDLR Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VLDLR Antibody (C-Term) - Images



All lanes : Anti-VLDLR Antibody (C-Term) at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 96 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

VLDLR Antibody (C-Term) - Background

Binds VLDL and transports it into cells by endocytosis. In order to be internalized, the receptor-ligand complexes must first cluster into clathrin-coated pits. Binding to Reelin induces tyrosine phosphorylation of Dab1 and modulation of Tau phosphorylation (By similarity).

VLDLR Antibody (C-Term) - References

Gafvels M.E., et al. Somat. Cell Mol. Genet. 19:557-569(1993).
Webb J.C., et al. Hum. Mol. Genet. 3:531-537(1994).
Sakai J., et al. J. Biol. Chem. 269:2173-2182(1994).
Oka K., et al. Genomics 20:298-300(1994).
Humphray S.J., et al. Nature 429:369-374(2004).